

GCE

Mathematics

Unit **4727**: Further Pure Mathematics 3

Advanced GCE

Mark Scheme for June 2014

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2014

Question	Answer	Marks	Guidance
1 (i)	$\begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \times \begin{pmatrix} 3 \\ 5 \\ 2 \end{pmatrix} = \begin{pmatrix} 7 \\ -7 \\ 7 \end{pmatrix} = 7 \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ <p>(eg) $z = 0 \Rightarrow 2x + y = 4, 3x + 5y = 13 \Rightarrow x = 1, y = 2$</p> $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$	<p>M1 A1</p> <p>M1</p> <p>A1</p>	<p>M1 requires evidence of method for cross product or at least 2 correct values calculated</p> <p>or any valid point e.g.(0, 3, -1), (3, 0, 2)</p> <p>Must have full equation including 'r ='</p> <p>oe vector form</p>
	<p>Alternative: Find one point Find a second point and vector between points</p> <p>multiple of $\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$</p> $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ <p>Alternative: Solve simultaneously</p> <p>Point and direction found</p> $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$	<p>M1 M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>M1 A1</p> <p>A1</p> <p>[4]</p>	<p>to at least expressions for x,y,z parametrically, or two relationship between 2 variables.</p>

Question	Answer	Marks	Guidance
1 (ii)	$\frac{ 2 \times 2 + 5 - -2 - 4 }{\sqrt{2^2 + 1^2 + 1^2}} = \frac{7}{\sqrt{6}}$	M1 A1	Condone lack of absolute signs for M1 2.86 with no workings scores M1 oe surd form. isw
	Alternative: find parameter for perpendicular meets plane and use to find distance	M1 [2]	For complete method with calculation errors look for $\lambda = -7/6$
2	$u = y^2 \Rightarrow \frac{du}{dx} = 2y \frac{dy}{dx}$ <p>so DE $\Rightarrow 2y \frac{dy}{dx} - 4y^2 = 2e^x$</p> $\Rightarrow \frac{du}{dx} - 4u = 2e^x$ $I = \exp \int -4 dx = e^{-4x}$ $e^{-4x} \frac{du}{dx} - 4e^{-4x} u = 2e^{-3x}$ $u e^{-4x} = -\frac{2}{3} e^{-3x} (+A)$ $u = -\frac{2}{3} e^x + A e^{4x}$ $y = \sqrt{-\frac{2}{3} e^x + A e^{4x}}$ <p>Alternative from 4th mark to 6th mark CF: $(u = \dots) A e^{4x}$</p> <p>PI: $u = k e^x, \frac{du}{dx} = k e^x$</p> $k e^x - 4k e^x = 2e^x, \quad k = -\frac{2}{3}$	M1 M1 A1 A1ft M1* *M1dep* M1dep* A1 A1 M1* M1 dep* [8]	Correctly finds or for complete unsimplified substitution Can be implied by next A1 Must have form $\frac{du}{dx} + f(x)u = g(x)$ for this mark and any further marks Can be implied by subsequent work No more than 1 numerical error at this step ignore use of '±'

Question	Answer	Marks	Guidance
3 (i)	$z^6 = 1 \Rightarrow z = e^{2k\pi i/6}$ $k = 0, 1, 2, 3, 4, 5$ Diagram	M1 A1 B1 B1 [4]	Oe exactly 6 roots 6 roots in right quadrant, correct angles and moduli accept roots 1, -1 given as integers. as evidenced by labels, circles, or accurate diagram, or by co-ordinates
3 (ii)	$(1+i)^6 = \left(\sqrt{2}e^{\frac{1}{4}\pi i}\right)^6$ $8e^{\frac{6}{4}\pi i}$ $= -8i$ <p>Alternative:</p> $(1+i)^6 = 1 + 6i + 15i^2 + 20i^3 + 15i^4 + 6i^5 + i^6$ $= 1 + 6i - 15 - 20i + 15 + 6i - 1$ $= -8i$ <p>Alternative: $(1+i)^2 = 2i$</p> $(1+i)^6 = (2i)^3$ $= -8i$	M1 M1 A1 M1 M1 A1 M1 M1 A1 [3]	Attempts modulus-argument form, getting at least 1 correct for $(\text{mod})^6$ and $\arg x$ ag complete argument including start line no more than 1 term wrong ag ag Sc 2 for only lines 2 & 3 correct

Question		Answer									Marks	Guidance																			
4	(i)										B1	2 or more	Ignore 1																		
		<table border="1"> <tr> <td>element</td> <td>(1)</td> <td>3</td> <td>7</td> <td>9</td> <td>11</td> <td>13</td> <td>17</td> <td>19</td> </tr> <tr> <td>inverse</td> <td>(1)</td> <td>7</td> <td>3</td> <td>9</td> <td>11</td> <td>17</td> <td>13</td> <td>19</td> </tr> </table>									element	(1)		3	7	9	11	13	17	19	inverse	(1)	7	3	9	11	17	13	19	B1	4 or more
		element	(1)	3	7	9	11	13	17	19																					
		inverse	(1)	7	3	9	11	17	13	19																					
									B1	all 7 correct																					
									[3]																						
4	(ii)	<p>(1 has order 1) 9,11,19 have order 2</p> <p>$3^2 = 9 \Rightarrow 3^4 = 1$ so order 4 similarly 7,13,17 order 4</p> <p>no element of order 8 so not cyclic</p>									M1	Correctly identifies order of all elements	Allow one error must show workings towards a^4 for demonstration that these elements are order 4 condone “no generator” in place of “no element of order 8”																		
											B1	justifies order for at least 1 element of order 4																			
											A1	www																			
											[3]																				
4	(iii)										M1	For two sets which both contain “1” and all (4) elements’ inverses																			
		<p>{1,13, 9, 17} and {1, 3, 9, 7}</p>									B1	One subgroup of order 4																			
											A1																				
											M1	for correspondence of “their” elements of same order																			
											A1	or $3 \leftrightarrow 17, 7 \leftrightarrow 13$																			
									[5]																						

Question	Answer	Marks	Guidance
5	<p>AE: $\lambda^2 + 5\lambda + 6 = 0$ $\lambda = -2, -3$ CF: $Ae^{-2x} + Be^{-3x}$ PI: $y = ae^{-x}$ $ae^{-x} - 5ae^{-x} + 6ae^{-x} = e^{-x}$ $2a = 1$ $a = \frac{1}{2}$ GS: $(y =) \frac{1}{2}e^{-x} + Ae^{-2x} + Be^{-3x}$ $x = 0, y = 0 \Rightarrow \frac{1}{2} + A + B = 0$ $y' = -\frac{1}{2}e^{-x} - 2Ae^{-2x} - 3Be^{-3x}$ $x = 0, y' = 0 \Rightarrow -\frac{1}{2} - 2A - 3B = 0$ $A = -1, B = \frac{1}{2}$ $y = \frac{1}{2}e^{-x} - e^{-2x} + \frac{1}{2}e^{-3x}$</p>	<p>B1 B1ft B1ft M1 A1 A1ft M1 M1* M1dep* A1 [10]</p>	<p>Differentiate and substitute Use condition on GS Differentiate their GS of form $y = ke^{-x} + Ae^{mx} + Be^{nx}$ where k, m, n are non-zero constants and m, n not 1 Use condition and attempt to find A, B www fit must be of form “ke^{-x} plus a standard CF form” with 2 arbitrary constants Must have 2 arbitrary constants Must have ‘y =’</p>

Question	Answer	Marks	Guidance
6 (i)	$l \parallel \begin{pmatrix} 2 \\ 3 \\ 5 \end{pmatrix} \quad \Pi \perp \begin{pmatrix} 4 \\ -1 \\ -1 \end{pmatrix} \quad \text{so} \quad \begin{pmatrix} 2 \\ 3 \\ 5 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ -1 \\ -1 \end{pmatrix} = 0 \Rightarrow l \parallel \Pi$ <p>$(1, -2, 7)$ on l but $4 \times 1 - 2 - 7 = -1 \neq 8$ so not in Π</p> <p>hence l not in Π</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>dot product of correct vectors = 0</p> <p>substitute point on line into Π and calculate d</p> <p>Full argument includes key components</p> <p>Argument can be about a general point on line</p>
6 (ii)	$(\mathbf{r} =) \begin{pmatrix} 1 \\ -2 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ -1 \end{pmatrix}$ <p>closest point where meets Π</p> $4(1 + 4\lambda) - (-2 - \lambda) - (7 - \lambda) = 8$ $\Rightarrow \lambda = \frac{1}{2}$ $\Rightarrow \mathbf{r} = \begin{pmatrix} 3 \\ -\frac{5}{2} \\ \frac{13}{2} \end{pmatrix}$	<p>B1</p> <p>M1</p> <p>A1ft</p> <p>A1</p> <p>[4]</p>	<p>parametric form of (x, y, z) substituted into plane</p>
6 (iii)	$\mathbf{r} = \begin{pmatrix} 3 \\ -\frac{5}{2} \\ \frac{13}{2} \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 3 \\ 5 \end{pmatrix}$	<p>B1ft</p> <p>[1]</p>	<p>oe</p> <p>must have "$\mathbf{r} =$"</p>

Question	Answer	Marks	Guidance
7 (i)	$2i \sin \theta = e^{i\theta} - e^{-i\theta}$ $2i \sin n\theta = e^{in\theta} - e^{-in\theta}$ $(2i \sin \theta)^5 = (e^{i\theta} - e^{-i\theta})^5$ $= e^{i5\theta} - 5e^{i3\theta} + 10e^{i\theta} - 10e^{-i\theta} + 5e^{-i3\theta} - e^{-i5\theta}$ $32i \sin^5 \theta = (e^{5i\theta} - e^{-5i\theta}) - 5(e^{3i\theta} - e^{-3i\theta}) + 10(e^{i\theta} - e^{-i\theta})$ $= 2i \sin 5\theta - 5(2i \sin 3\theta) + 10(2i \sin \theta)$ $\sin^5 \theta = \frac{1}{16}(\sin 5\theta - 5 \sin 3\theta + 10 \sin \theta)$	B1 M1* M1dep* A1 [4]	any equivalent form binomial expansion grouping terms AG If use z , must define it can be unsimplified Award B1 then sc M1A1 for candidates who omit this stage from otherwise complete argument must convince on the $\frac{1}{16}$ and on the elimination of i
7 (ii)	$16 \sin^5 \theta - 10 \sin \theta = \sin 5\theta - 5 \sin 3\theta$ $16 \sin^5 \theta - 6 \sin \theta = 0$ $\sin \theta = 0, \pm \sqrt[4]{\frac{3}{8}}$ $\theta = 0, \pm 0.899$	M1* A1 M1dep* A1 [4]	Attempts to eliminate $\sin 5\theta$ and $\sin 3\theta$ must have 3 values for $\sin \theta$ Or $16 \sin^5 \theta = 6 \sin \theta$

Question	Answer	Marks	Guidance
8 (i)	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ is identity}$ $\begin{pmatrix} a & -b \\ b & a \end{pmatrix}^{-1} = \frac{1}{a^2 + b^2} \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \in G$ $\begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} c & -d \\ d & c \end{pmatrix} = \begin{pmatrix} ac - bd & -bc - ad \\ bc + ad & ac - bd \end{pmatrix}$ <p>and</p> $(ac - bd)^2 + (bc + ad)^2 = a^2c^2 + b^2d^2 + b^2c^2 + a^2d^2$ $= (a^2 + b^2)(c^2 + d^2) \neq 0$	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>for M1, must at least get matrix in form $\begin{pmatrix} x & -y \\ y & x \end{pmatrix}$, or state existence of inverse due to non-singularity</p> <p>Must not attempt to prove commutativity in (i)</p>
8 (ii)	$\begin{pmatrix} c & -d \\ d & c \end{pmatrix} \begin{pmatrix} a & -b \\ b & a \end{pmatrix} = \begin{pmatrix} ac - bd & -bc - ad \\ bc + ad & ac - bd \end{pmatrix}$ $= \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} c & -d \\ d & c \end{pmatrix} \text{ so commutative}$	<p>M1</p> <p>A1</p> <p>[2]</p>	<p>must also consider matrices reversed, but may be seen in (i)</p>
8 (iii)	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^2 = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}^2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ <p>order 4</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>g^2 must be correct</p> <p>allow 1 error in getting g^4</p>

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2014

