

GCE

Mathematics

Unit 4730: Mechanics 3

Advanced GCE

Mark Scheme for June 2017

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

Answer		Marks	Guidance		
1	(i)	Impulse/momentum triangle with sides 0.8, 1.2 and 1 $\cos \theta = \frac{0.8^2 + 1^2 - 1.2^2}{2 \times 0.8 \times 1}$ 82.8° or 1.44 rads 1.2 sin $\alpha = \sin \theta$ Angle 124°	B1 M1 A1 M1 A1 [5]	OR $1.2 \cos \alpha = \cos \theta - 0.8$ $1.2 \sin \alpha = \sin \theta$ M1 $1.44 = (\cos \theta - 0.8)^2 + \sin^2 \theta$ A1 isw cv θ ; OR from cos rule No isw	Square and add 82.81924° or 1.445 rads may see 55.771° or 0.97339 rads 2.168 rads
2	(i)	$\frac{1}{2}m \times 0.7^2 = \frac{1}{2}mv^2 + \frac{24mg0.3^2}{2 \times 1.2} - mg \times 1.5$ Speed = 3.5 (ms ⁻¹)	M1 A1 A1 [3]	By energy; needs KE, PE and EE terms $OR \frac{1}{2}m \times 4.9^2 = \frac{1}{2}mv^2 + \frac{24mg0.3^2}{2\times 1.2} - mg \times 0.3$ AG Adequate working, no errors	Allow wrong signs, missing '2'
	(ii)	One correct EE term involving x seen $\frac{1}{2}m \times 0.7^{2} = \frac{24mg(x-1.2)^{2}}{2 \times 1.2} + \frac{32mg(x-1.5)^{2}}{2 \times 0.8} - mgx$ [48x ² - 136x + 95 = 0] 1.25 (m) and 1.58 (m)	B1 M1 A1 M1 A1 [5]	Where x is distance below O OR, where x is dist from T, $\frac{1}{2}m \times 0.7^2 = \frac{24mg(x+0.3)^2}{2\times 1.2} + \frac{32mgx^2}{2\times 0.8} - mg(x+1.5)$ Leads to $48x^2 + 8x - 1 = 0$ Correct attempt to solve their 3 term quad. $1\frac{1}{4} + 1\frac{7}{12}$	Energy equation with at least 1 KE, 1 PE and 1 EE term and values subst. Alt left side: $\frac{1}{2}m3.5^2 + \frac{24mg0.3^2}{2\times1.2} - 1.5mg$ Dep M1 above

Mark Scheme

3	(i)	$48 (ms^{-1})$	B1 [1]	Accept \leq 48	
	(ii)	Use $\frac{1}{2}\sqrt{12 - \frac{1}{4}v} = 0.2a$	M1*	Accept $v \frac{\mathrm{d}v}{\mathrm{d}x}$ for a	Allow missing 0.2 or sign error
		$\frac{1}{2} \sqrt{12 - \frac{1}{4}v} = 0.2 \frac{dv}{dt}$	A1		
		$2.5t = \int \frac{dv}{\sqrt{12 - \frac{1}{2}v}} (+c)$	*M1*	Sep variables and integrate one side	$2.5t = \int \frac{2\mathrm{d}v}{\sqrt{48-v}} (+c)$
		$2.5t = -8\left(12 - \frac{1}{4}v\right)^{\frac{1}{2}}(+c)$	A1		$2.5t = -4(48 - v)^{\frac{1}{2}}(+c)$
		[c = 24] v = 48 - 4 $\left(3 - \frac{t}{3.2}\right)^2$	*M1 A1 [6]	For attempt to find c, dep previous M1 oe $12 + 7.5t - \frac{25}{64}t^2$ (0.390625)	$v = 48 - 0.390625(9.6 - t)^2$
	(iii)	$x = \int (12 + \frac{24}{3.2}t - \frac{4}{3.2^2}t^2) \mathrm{d}t$	M1	OR $x = \int (48 - 4\left(3 - \frac{t}{3.2}\right)^2) dt$	OR $\frac{1}{2}\sqrt{12 - \frac{1}{4}v} = 0.2v\frac{dv}{dx}$
		$x = 12t + 3.75t^2 - 0.1302t^3(+c)$	A1	$x = 48t + \frac{12.8}{3} \left(3 - \frac{t}{32}\right)^3 (+c')$	via subst $\left(12 - \frac{1}{4}v\right) = u^2$
		(t = 0 and) t = 3.2	M1	ft their (ii)	$x = 12.8\left(12u - \frac{u^3}{3}\right) + C$
		Distance = 72.533 (m)	A1 [4]		×

4	(i)	Momentum equation $2ma = -2m\frac{1}{10}\sqrt{5}\frac{1}{\sqrt{5}} + 7m\frac{1}{10}\sqrt{5}\frac{1}{\sqrt{5}}$	M1	Along line of centres Allow errors with signs and masses	Allow use of 63.4° for full marks Must use comp of vel
		$10 \sqrt{5} 10 \sqrt{5}$ $(a =) \frac{1}{4} (\text{ms}^{-1})$ Comp of speed of A perp = 0.2 Speed of A was $\sqrt{(0.25^2 + 0.2^2)}$ OR tan $\theta = \frac{0.2}{0.25}$ Speed 0.320 or $\frac{\sqrt{41}}{20}$; Ang 38.7° or 0.675 rads NLM 0.1 + 0.1 = $-e(0 - a)$) (e =) 0.8	A1 B1 M1 A1 M1 A1 [7]	soi soi Allow their vel comps oe For both angle and speed Along line of centres Allow errors with signs	0.320156; 38.6598° or 0.67474 rads May see $\frac{1}{10}\sqrt{5}\frac{1}{\sqrt{5}}$ for 0.1
	(ii)	A and B have same speed perpendicular to line of centres after first collision	B1 [1]	accept 'vertical'	
	(iii)	Momentum equation along line of centres $3mU - 2m\frac{1}{10}\sqrt{5}\frac{1}{\sqrt{5}} = 3mc + 2ma'$ NLM $a' - c = -1(-\frac{1}{10}\sqrt{5}\frac{1}{\sqrt{5}} - U)$ Use $a' = 0.1$ Max $U = \frac{1}{15}$	M1 A1 M1 A1 B1 A1 [6]	Allow errors with signs and masses Allow $\cos \alpha$ for $\frac{1}{\sqrt{5}}$ Allow errors with signs Accept any inequality Accept 0.0667 accept \leq	Must use comp of vel Or conservation of energy $\frac{1}{2}3mU^2 + \frac{1}{2}2m0.1^2 = \frac{1}{2}3mc^2 + \frac{1}{2}2ma'^2$ do not accept <
5	(i)	$3mga\cos\frac{\pi}{6} \text{ and } 2mga\cos\frac{\pi}{6}$ $3mga\cos\left(\frac{\pi}{6} + \theta\right) + 2mga\cos\left(\frac{\pi}{6} - \theta\right) + \frac{1}{2}3mv^{2} + \frac{1}{2}2mv^{2}$ $v^{2} = \frac{2}{5}ag\left(5\cos\frac{\pi}{6} - 3\cos\left(\frac{\pi}{6} + \theta\right) - 2\cos\left(\frac{\pi}{6} - \theta\right)\right)$	B1 M1 A1 A1 [4]	Initial PE Final PE + KE AG Equating and correct manipulation	If <i>O</i> is zero level for PE For M1 at least 1 KE and 1 PE term; allow m used for 2m/3m; wrong signs; missing <i>g</i>

	(ii)	$v^{2} = \frac{2}{5}ag\left(5\cos\frac{\pi}{6} - 3\cos\frac{\pi}{3} - 2\cos 0\right)$ $3mg\cos\frac{\pi}{3} - R = 3m\frac{v^{2}}{a}$ $R = 3mg\cos\frac{\pi}{3} - 3m\frac{2g}{5}\left(5\cos\frac{\pi}{6} - \frac{7}{2}\right)$	B1 M1 A1	$v^2 = \frac{1}{5}ag(5\sqrt{3} - 7)$ F = ma, condone sign error; allow m used for 2m/3m	OR $3mg \cos\left(\frac{\pi}{6} + \theta\right) - R = 3m\frac{v^2}{a}$ $R = 3mg \cos\left(\frac{\pi}{6} + \theta\right) - 3m\frac{2}{5}g\left(5\cos\frac{\pi}{6} - 3\cos\left(\frac{\pi}{6} + \theta\right) - 2\cos\left(\frac{\pi}{6} - \theta\right)\right)$ $R = mg(6.6\cos\left(\frac{\pi}{6} + \theta\right) + 2.4\cos\left(\frac{\pi}{6} - \theta\right) - 6\cos\frac{\pi}{6})$
		$R = mg\left(5.7 - 6\cos\frac{\pi}{6}\right)$ oe	A1 [4]	Accept 0.5038475 <i>mg</i> or $mg(5.7 - 3\sqrt{3})$ oe	Answer must be simplified 4.94 <i>m</i> loses last mark
6	(i)	$Pl\sqrt{5} = W \times 3l \cos \theta$ P = 1.2W $Ql\sqrt{2} = U \times \frac{\lambda}{2}l \cos \phi$ $O = 0.25\lambda U$	M1 A1 M1 A1 [4]	Mom about <i>A</i> for <i>AB</i> AG Mom about A for AC	Allow sin θ , cancelled <i>l</i> Not from use of angle 26.565° Allow sin \emptyset , cancelled <i>l</i>
	(ii)	(H) $P \sin \theta = Q \sin \phi$ (V) $W + U = P \cos \theta + Q \cos \phi$ $W + U = P \cos \theta + P \sin \theta \times \frac{\cos \phi}{\sin \phi}$ $W + U = \frac{3}{\sqrt{5}} \times 1.2W$ k = 0.610 $\lambda = 4.98$ $[P\sqrt{5}l - W3l \cos \theta = Q\sqrt{2}l - U\frac{\lambda}{2}\cos \phi]$	M1 M1 A1 M1* *M1 A1 A1 [7]	$P\frac{1}{\sqrt{5}} = Q\frac{1}{\sqrt{2}}; \text{ compts essential}$ $W + U = P\frac{2}{\sqrt{5}} + Q\frac{1}{\sqrt{2}}; \text{ compts essential}$ Eliminate Q (or P) dep M1M1 Elim P and Q to get equation in k, W + U = 1.609689W $\frac{18\sqrt{5} - 25}{25}$ Mom about A (or any other point) for whole system – allow M1(A1) if resolving not seen twice]	Allow $\frac{2}{\sqrt{5}}$ for M1 Allow $\frac{1}{\sqrt{5}}$ for M1, sign errors $W + U = Q \cos \theta \times \frac{\sin \phi}{\sin \theta} + Q \cos \phi$ $\left[W + U = 0.25\lambda U \times \frac{3}{2\sqrt{2}}\right]$ 0.6099689 4.97695 Allow use of angles in (ii): 26.6 & 45 OR after M1M1A0/1, M1* for 2 equns in terms of k and λ , *M1 for solving for k or λ .

7	(i)	$\frac{1}{2}m \times \frac{g}{90} = mgh$ [Max height = $\frac{1}{180}$ = 0.005556]	M1	By energy; allow cancelled m	$\frac{1}{2}m \times \frac{g}{90} = mg \times 0.8(1 - \cos\theta)$
		Max angle = 6.76° or 0.118 rads	AI M1	Allow MIAI for 6.76° or 0.118 rads in (ii)	0.75070.11798
		$-mg\sin\theta = m \times 0.8 \times \theta$		N2L; allow <i>a</i> for 0.8 θ ; allow cancelled <i>m</i>	anow sign error, sin / cos
		$\theta = -\frac{1}{0.8}\theta,$	ЛІ		
		SHM (about $\theta = 0$) since θ is small	Al		
		$\omega^2 = 12.25$		Cand value	1 7052
		Period = 1.80 secs $(\frac{1}{7}\pi)$	AI [/]		1.7952
	(ii)	$0.087266 = A \sin 3.5t$ $t = 0.238 \sec t$ $t' = 2\left(\frac{1.7952}{4} - 0.2378\right) = 0.422(s)$	M1 A1 A1	OR $5 = A \sin 3.5t$; $A =$ amplitude Or 0.65972 Or 0.65972 – 0.2378	May use cos * 0.2378
		$\dot{\theta} = 0.118 \times 3.5 \cos 3.5 \times 0.238$	M1	OR $\dot{\theta} = \sqrt{(3.5^2(0.118^2 - 0.0873^2))}$	allow sin if consistent with $*$; allow 5° and 6.76°
		Linear speed = $0.222(ms^{-1})$	A1 [5]	0.8 ×0.278	Or $\frac{1}{2}m\frac{9}{90} = \frac{1}{2}mv^2 + mg0.8(1 - \cos 5^\circ)$
	(iii)	Max height is still 0.00556 soi	B1	accept 'still the same'	or attempt to work out height
		Max angle = $\cos^{-1} \frac{(0.05 - 0.00556)}{0.05}$ [27.3]	B1		0.476 rads
		Not SHM since angle is not small	B1 [3]		

OCR (Oxford Cambridge and RSA Examinations) 1 Hills Road Cambridge CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998 Facsimile: 01223 552627 Email: <u>general.qualifications@ocr.org.uk</u>

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee Registered in England Registered Office; 1 Hills Road, Cambridge, CB1 2EU Registered Company Number: 3484466 OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations) Head office Telephone: 01223 552552 Facsimile: 01223 552553 PART OF THE CAMBRIDGE ASSESSMENT GROUP

