

# Mark Scheme (Final) Summer 2007

GCE

GCE Mathematics (6665/01)

**June 2007**  
**6665 Core Mathematics C3**  
**Mark Scheme**

Question Number	Scheme	Marks
<b>1.</b>	<p>(a) <math>\ln 3x = \ln 6</math> or <math>\ln x = \ln \left(\frac{6}{3}\right)</math> or <math>\ln \left(\frac{3x}{6}\right) = 0</math>  <math>x = 2</math> (only this answer)</p>	<p>M1  A1 (cso) (2)</p>
	<p>(b) <math>(e^x)^2 - 4e^x + 3 = 0</math> (any 3 term form)  <math>(e^x - 3)(e^x - 1) = 0</math>  <math>e^x = 3</math> or <math>e^x = 1</math> Solving quadratic  <math>x = \ln 3,</math> <math>x = 0</math> (or <math>\ln 1</math>)</p>	<p>M1  M1 dep  M1 A1 (4)  <b>(6 marks)</b></p>

Notes: (a) Answer  $x = 2$  with no working or no incorrect working seen: M1A1

Note:  $x = 2$  from  $\ln x = \frac{\ln 6}{\ln 3} = \ln 2$  M0A0

$\ln x = \ln 6 - \ln 3 \Rightarrow x = e^{(\ln 6 - \ln 3)}$  allow M1,  $x = 2$  (no wrong working) A1

- (b) 1<sup>st</sup> M1 for attempting to multiply through by  $e^x$ : Allow  $y, X$ , even  $x$ , for  $e^x$   
2<sup>nd</sup> M1 is for solving quadratic as far as getting two values for  $e^x$  or  $y$  or  $X$  etc  
3<sup>rd</sup> M1 is for converting their answer(s) of the form  $e^x = k$  to  $x = \ln k$  (must be exact)  
A1 is for  $\ln 3$  **and**  $\ln 1$  or 0 (Both required and no further solutions)

2. (a)	$2x^2 + 3x - 2 = (2x - 1)(x + 2)$ $f(x) = \frac{(2x+3)(2x-1) - (9+2x)}{(2x-1)(x+2)}$ at any stage f.t. on error in denominator factors (need not be single fraction) Simplifying numerator to quadratic form  Correct <b>numerator</b> $= \frac{4x^2 + 2x - 12}{[(2x-1)(x+2)]}$ Factorising numerator, <b>with</b> a denominator $= \frac{2(2x-3)(x+2)}{(2x-1)(x+2)}$ o.e. $= \frac{4x-6}{2x-1}$ (*)	B1 M1, A1√ M1 A1 M1 A1 cso (7)
Alt.(a)	$2x^2 + 3x - 2 = (2x - 1)(x + 2)$ $f(x) = \frac{(2x+3)(2x^2 + 3x - 2) - (9+2x)(x+2)}{(x+2)(2x^2 + 3x - 2)}$ $= \frac{4x^3 + 10x^2 - 8x - 24}{(x+2)(2x^2 + 3x - 2)}$ $= \frac{2(x+2)(2x^2 + x - 6)}{(x+2)(2x^2 + 3x - 2)}$ at any stage B1 M1A1 f.t. o.e. or $\frac{2(2x-3)(x^2 + 4x + 4)}{(x+2)(2x^2 + 3x + 2)}$ o.e. Any one linear factor × quadratic factor in <b>numerator</b> M1, A1 $= \frac{2(x+2)(x+2)(2x-3)}{(x+2)(2x^2 + 3x - 2)}$ o.e. M1 $= \frac{2(2x-3)}{2x-1} \quad \frac{4x-6}{2x-1}$ (*) A1	
(b)	Complete method for $f'(x)$ ; e.g. $f'(x) = \frac{(2x-1) \times 4 - (4x-6) \times 2}{(2x-1)^2}$ o.e. $= \frac{8}{(2x-1)^2}$ or $8(2x-1)^{-2}$ Not treating $f^{-1}$ (for $f'$ ) as misread	M1 A1 A1 (3) <b>(10 marks)</b>

Notes: (a) 1<sup>st</sup> M1 in either version is for correct method

1<sup>st</sup> A1 Allow  $\frac{2x+3(2x-1) - (9+2x)}{(2x-1)(x+2)}$  or  $\frac{(2x+3)(2x-1) - 9 + 2x}{(2x-1)(x+2)}$  or  $\frac{2x+3(2x-1) - 9 + 2x}{(2x-1)(x+2)}$  (fractions)

2<sup>nd</sup> M1 in (main a) is for forming 3 term quadratic in **numerator**

3<sup>rd</sup> M1 is for factorising their quadratic (usual rules) ; factor of 2 need not be extracted

(\*) A1 is given answer so is cso


Alt :(a) 3<sup>rd</sup> M1 is for factorising resulting quadratic

(b) SC: For M allow  $\pm$  given expression or one error in product rule

Alt: Attempt at  $f(x) = 2 - 4(2x-1)^{-1}$  and diff. M1;  $k(2x-1)^{-2}$  A1; A1 as above

Accept  $8(4x^2 - 4x + 1)^{-1}$ .

Differentiating original function – mark as scheme.

Question Number	Scheme	Marks
3. (a)	$\frac{dy}{dx} = x^2e^x + 2xe^x$	M1,A1,A1 (3)
(b)	If $\frac{dy}{dx} = 0$ , $e^x(x^2 + 2x) = 0$ setting $(a) = 0$ $[e^x \neq 0]$ $x(x + 2) = 0$ $(x = 0)$ $x = -2$ $x = 0, y = 0$ <b>and</b> $x = -2, y = 4e^{-2} (= 0.54\dots)$	M1  A1 A1 $\checkmark$ (3)
(c)	$\frac{d^2y}{dx^2} = x^2e^x + 2xe^x + 2xe^x + 2e^x$ $[= (x^2 + 4x + 2)e^x]$	M1, A1 (2)
(d)	$x = 0, \frac{d^2y}{dx^2} > 0 (=2)$ $x = -2, \frac{d^2y}{dx^2} < 0 [= -2e^{-2} (= -0.270\dots)]$ M1: Evaluate, or state sign of, candidate's (c) for at least one of candidate's x value(s) from (b) $\therefore$ minimum $\therefore$ maximum	M1  A1 (cso) (2)
Alt.(d)	For M1: Evaluate, or state sign of, $\frac{dy}{dx}$ at two appropriate values – on either side of at least one of their answers from (b) or Evaluate y at two appropriate values – on either side of at least one of their answers from (b) or Sketch curve 	

**(10 marks)**

Notes: (a) M for attempt at  $f(x)g'(x) + f'(x)g(x)$

1<sup>st</sup> A1 for one correct, 2<sup>nd</sup> A1 for the other correct.

**Note that  $x^2e^x$  on its own scores no marks**

(b) 1<sup>st</sup> A1 ( $x = 0$ ) may be omitted, but for

2<sup>nd</sup> A1 both sets of coordinates needed ; f.t only on candidate's  $x = -2$


(c) M1 requires complete method for candidate's (a), result may be unsimplified for A1

(d) A1 is cso;  $x = 0$ , min, and  $x = -2$ , max and no incorrect working seen,

or (in alternative) sign of  $\frac{dy}{dx}$  either side correct, or values of y appropriate to t.p.

Need only consider the quadratic, as may assume  $e^x > 0$ .

**If all marks gained** in (a) and (c), and correct x values, give M1A1 for correct statements with no working

Question Number	Scheme	Marks	
4.	(a) $x^2(3-x) - 1 = 0$ o.e. (e.g. $x^2(-x+3) = 1$ ) $x = \sqrt{\frac{1}{3-x}}$ (*) Note(*), answer is given: need to see appropriate working and A1 is cso [Reverse process: Squaring and non-fractional equation M1, form f(x) A1]	M1 A1 (cso) (2)	
	(b) $x_2 = 0.6455, x_3 = 0.6517, x_4 = 0.6526$ 1 <sup>st</sup> B1 is for one correct, 2 <sup>nd</sup> B1 for other two correct If all three are to greater accuracy, award B0 B1	B1; B1 (2)	
	(c) Choose values in interval (0.6525, 0.6535) or tighter and evaluate both $f(0.6525) = -0.0005$ (372... $f(0.6535) = 0.002$ (101... At least one correct "up to bracket", i.e. -0.0005 or 0.002 <b>Change of sign</b> , $\therefore x = 0.653$ is a root (correct) to 3 d.p. Requires both correct "up to bracket" and conclusion as above	M1 A1 A1 (3) <b>(7 marks)</b>	
	Alt (i) Continued iterations at least as far as $x_6$ M1 $x_5 = 0.65268, x_6 = 0.6527, x_7 = \dots$ two correct to at least 4 s.f. A1 Conclusion : Two values correct to 4 d.p., so 0.653 is root to 3 d.p. A1 Alt (ii) If use $g(0.6525) = 0.6527.. > 0.6525$ and $g(0.6535) = 0.6528.. < 0.6535$ M1A1 Conclusion : Both results correct, so 0.653 is root to 3 d.p. A1		
5.	(a) Finding $g(4) = k$ and $f(k) = \dots$ or $fg(x) = \ln\left(\frac{4}{x-3} - 1\right)$ $[f(2) = \ln(2 \times 2 - 1) \quad fg(4) = \ln(4 - 1)] = \ln 3$	M1 A1 (2)	
	(b) $y = \ln(2x-1) \Rightarrow e^y = 2x-1$ or $e^x = 2y-1$ $f^{-1}(x) = \frac{1}{2}(e^x + 1)$ Allow $y = \frac{1}{2}(e^x + 1)$ Domain $x \in \mathcal{R}$ [Allow $\mathcal{R}$ , all reals, $(-\infty, \infty)$ ] independent	M1, A1 A1 B1 (4)	
	(c) 	Shape, and x-axis should appear to be asymptote <b>Equation x = 3 needed</b> , may see in diagram (ignore others) Intercept $(0, \frac{2}{3})$ no other; accept $y = \frac{2}{3}$ (0.67) or on graph	B1 B1 ind. B1 ind (3)
	(d) $\frac{2}{x-3} = 3 \Rightarrow x = 3\frac{2}{3}$ or exact equiv. $\frac{2}{x-3} = -3, \Rightarrow x = 2\frac{1}{3}$ or exact equiv. Note: $2 = 3(x+3)$ or $2 = 3(-x-3)$ o.e. is M0A0 Alt: Squaring to quadratic $(9x^2 - 54x + 77 = 0)$ and solving M1; B1A1	B1 M1, A1 (3) <b>(12 marks)</b>	

6.	(a)	Complete method for $R$ : e.g. $R \cos \alpha = 3$ , $R \sin \alpha = 2$ , $R = \sqrt{3^2 + 2^2}$ $R = \sqrt{13}$ or 3.61 (or more accurate) Complete method for $\tan \alpha = \frac{2}{3}$ [Allow $\tan \alpha = \frac{3}{2}$ ] $\alpha = 0.588$ (Allow $33.7^\circ$ )	M1 A1 M1 A1 (4)
	(b)	Greatest value = $(\sqrt{13})^4 = 169$	M1, A1 (2)
	(c)	$\sin(x + 0.588) = \frac{1}{\sqrt{13}}$ (= 0.27735...) $\sin(x + \text{their } \alpha) = \frac{1}{\text{their } R}$ $(x + 0.588) = 0.281(03\dots)$ or $16.1^\circ$ $(x + 0.588) = \pi - 0.28103\dots$ Must be $\pi - \text{their } 0.281$ or $180^\circ - \text{their } 16.1^\circ$ or $(x + 0.588) = 2\pi + 0.28103\dots$ Must be $2\pi + \text{their } 0.281$ or $360^\circ + \text{their } 16.1^\circ$ $x = 2.273$ or $x = 5.976$ (awrt) Both (radians only) If 0.281 or $16.1^\circ$ not seen, correct answers imply this A mark	M1 A1 M1 M1 A1 (5) <b>(11 marks)</b>


Notes: (a) 1<sup>st</sup> M1 for correct method for  $R$   
 2<sup>nd</sup> M1 for correct method for  $\tan \alpha$   
 No working at all: M1A1 for  $\sqrt{13}$ , M1A1 for 0.588 or  $33.7^\circ$ .  
 N.B.  $R \cos \alpha = 2$ ,  $R \sin \alpha = 3$  used, can still score M1A1 for  $R$ , but loses the A mark for  $\alpha$ .  
 $\cos \alpha = 3$ ,  $\sin \alpha = 2$ : apply the same marking.

(b) M1 for realising  $\sin(x + \alpha) = \pm 1$ , so finding  $R^4$ .

(c) Working in mixed degrees/rads : first two marks available  
 Working consistently in degrees: Possible to score first 4 marks  
 [Degree answers, just for reference only, are  $130.2^\circ$  and  $342.4^\circ$ ]  
 Third M1 can be gained for candidate's  $0.281 - \text{candidate's } 0.588 + 2\pi$  or equiv. in degrees  
 One of the answers correct in radians or degrees implies the corresponding M mark.

Alt: (c) (i) Squaring to form quadratic in  $\sin x$  or  $\cos x$  M1  
 $[13 \cos^2 x - 4 \cos x - 8 = 0, 13 \sin^2 x - 6 \sin x - 3 = 0]$   
 Correct values for  $\cos x = 0.953\dots, -0.646$ ; or  $\sin x = 0.767, 2.27$  awrt A1  
 For any one value of  $\cos x$  or  $\sin x$ , correct method for two values of  $x$  M1  
 $x = 2.273$  or  $x = 5.976$  (awrt) Both seen anywhere A1  
 Checking other values (0.307, 4.011 or 0.869, 3.449) and discarding M1

(ii) Squaring and forming equation of form  $a \cos 2x + b \sin 2x = c$   
 $9 \sin^2 x + 4 \cos^2 x + 12 \sin 2x = 1 \Rightarrow 12 \sin 2x + 5 \cos 2x = 11$   
 Setting up to solve using R formula e.g.  $\sqrt{13} \cos(2x - 1.176) = 11$  M1  
 $(2x - 1.176) = \cos^{-1}\left(\frac{11}{\sqrt{13}}\right) = 0.562(0\dots)$  ( $\alpha$ ) A1  
 $(2x - 1.176) = 2\pi - \alpha, 2\pi + \alpha, \dots$  M1  
 $x = 2.273$  or  $x = 5.976$  (awrt) Both seen anywhere A1  
 Checking other values and discarding M1

Question Number	Scheme	Marks
<p>7. (a)</p> <p>Alt.(a)</p>	$\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = \frac{\sin^2 \theta + \cos^2 \theta}{\cos \theta \sin \theta}$ <p>M1 Use of common denominator to obtain single fraction</p> $= \frac{1}{\cos \theta \sin \theta}$ <p>M1 Use of appropriate trig identity (in this case <math>\sin^2 \theta + \cos^2 \theta = 1</math>)</p> $= \frac{1}{\frac{1}{2} \sin 2\theta}$ <p>Use of <math>\sin 2\theta = 2 \sin \theta \cos \theta</math></p> $= 2 \operatorname{cosec} 2\theta \quad (*)$ $\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = \tan \theta + \frac{1}{\tan \theta} = \frac{\tan^2 \theta + 1}{\tan \theta}$ <p>M1</p> $= \frac{\sec^2 \theta}{\tan \theta}$ <p>M1</p> $= \frac{1}{\cos \theta \sin \theta} = \frac{1}{\frac{1}{2} \sin 2\theta}$ <p>M1</p> $= 2 \operatorname{cosec} 2\theta \quad (*) \quad (\text{cso}) \quad \text{A1}$ <p>If show two expressions are equal, need conclusion such as QED, tick, true.</p>	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1 cso (4)</p>
(b)	 <p>Shape (May be translated but need to see 4 "sections")</p> <p>T.P.s at <math>y = \pm 2</math>, asymptotic at correct <math>x</math>-values (dotted lines not required)</p>	<p>B1</p> <p>B1 dep. (2)</p>
(c)	$2 \operatorname{cosec} 2\theta = 3$ $\sin 2\theta = \frac{2}{3} \quad \text{Allow } \frac{2}{\sin 2\theta} = 3 \quad [\text{M1 for equation in } \sin 2\theta]$ $(2\theta) = [41.810\dots^\circ, 138.189\dots^\circ; 401.810\dots^\circ, 498.189\dots^\circ]$ <p>1st M1 for <math>\alpha, 180 - \alpha</math>; 2nd M1 adding <math>360^\circ</math> to at least one of values</p> $\theta = 20.9^\circ, 69.1^\circ, 200.9^\circ, 249.1^\circ \quad (1 \text{ d.p.}) \quad \text{awrt}$	<p>M1, A1</p> <p>M1; M1</p>
<p>Note</p> <p>Alt.(c)</p>	<p>1st A1 for any two correct, 2nd A1 for other two Extra solutions in range lose final A1 only SC: Final 4 marks: <math>\theta = 20.9^\circ</math>, after M0M0 is B1; record as M0M0A1A0</p> $\tan \theta + \frac{1}{\tan \theta} = 3 \quad \text{and form quadratic, } \tan^2 \theta - 3 \tan \theta + 1 = 0 \quad \text{M1, A1}$ <p>(M1 for attempt to multiply through by <math>\tan \theta</math>, A1 for correct equation above)</p> <p>Solving quadratic <math>[\tan \theta = \frac{3 \pm \sqrt{5}}{2} = 2.618\dots \text{ or } = 0.3819\dots]</math> M1</p> $\theta = 69.1^\circ, 249.1^\circ \quad \theta = 20.9^\circ, 200.9^\circ \quad (1 \text{ d.p.}) \quad \text{M1, A1, A1}$ <p>(M1 is for one use of <math>180^\circ + \alpha^\circ</math>, A1A1 as for main scheme)</p>	<p>A1, A1 (6)</p> <p>(12 marks)</p>

Question Number	Scheme	Marks
8. (a)	$D = 10, t = 5, \quad x = 10e^{-\frac{1}{8} \times 5}$ $= 5.353$ awrt	M1 A1 (2)
(b)	$D = 10 + 10e^{-\frac{5}{8}}, t = 1, \quad x = 15.3526... \times e^{-\frac{1}{8}}$ $x = 13.549$ (*)	M1 A1 cso (2)
Alt.(b)	$x = 10e^{-\frac{1}{8} \times 6} + 10e^{-\frac{1}{8} \times 1}$ M1 $x = 13.549$ (*) A1 cso	
(c)	$15.3526...e^{-\frac{1}{8}T} = 3$ $e^{-\frac{1}{8}T} = \frac{3}{15.3526...} = 0.1954...$ $-\frac{1}{8}T = \ln 0.1954...$ $T = 13.06... \text{ or } 13.1 \text{ or } 13$	M1  M1  A1 (3)
		<b>(7 marks)</b>

Notes: (b) (main scheme) M1 is for  $(10 + 10e^{-\frac{5}{8}})e^{-\frac{1}{8}}$ , or  $\{10 + \text{their(a)}\}e^{-\frac{1}{8}}$

**N.B.** The answer is given. There are many correct answers seen which deserve M0A0  
or M1A0

(c) 1<sup>st</sup> M is for  $(10 + 10e^{-\frac{5}{8}})e^{-\frac{T}{8}} = 3$  o.e.

2<sup>nd</sup> M is for converting  $e^{-\frac{T}{8}} = k$  ( $k > 0$ ) to  $-\frac{T}{8} = \ln k$ . This is independent of 1<sup>st</sup> M.

Trial and improvement: M1 as scheme,  
M1 correct process for their equation (two equal to 3 s.f.)  
A1 as scheme