Question	Scheme	Marks	AOs
1(a)	$[Q_2 =] (5+) \frac{12}{15} \times 5$ or (use of $(n+1)$) $(5+) \frac{12.5}{15} \times 5$	M1	1.1a
	= 9 or 9.166 awrt 9.17	A1	1.1b
		(2)	
(b)	$\left[\sigma_x = \right] \sqrt{\frac{5675}{30} - \left(\frac{355}{30}\right)^2} = \sqrt{49.14}$	M1	1.1a
	= <u>awrt 7.01</u>	A1	1.1b
	Accept $s_x = \sqrt{\frac{5675 - 30\left(\frac{355}{30}\right)^2}{29}} = 7.1294$		
		(2)	
(c)	$x = \frac{t-15}{2}$ or $t = 2x+15$	M1	3.1b
	Median = 2' "9"+ 15 = 33 (allow awrt 33.3 from "9.17" in (a))	A1ft	1.1b
	Sd = 2' "7.01" = 14.02 (awrt 14.0) [allow awrt 14.3 if s used]	A1ft	1.1b
		(3)	
(d)	The median time is "33" and "33" < 35 so 50% (30) should finish in 35 minutes. ALT Probability of being < 35 mins is $\frac{18}{30} \setminus \frac{18}{30}$, $60 = 36$ applicants to choose from.	M1	2.4
	It is likely that they will fill all 25 positions [providing those offered accept]	A1	2.2b
		(2)	

Notes: (9 marks)

(a) M1: For a suitable fraction × 5 (ignore end points)

A1: For 9 or awrt 9.17 if using n + 1

(b) M1: For a correct expression for \overline{x} and s_x or s_x

A1: For awrt $s_x = 7.01$ or $s_x =$ awrt 7.13

(c) M1: For realising $x = \frac{t-15}{2}$ and then rearranging to get a correct equation with t as the subject

May be implied by a correct answer for the median of t.

A1ft: ft their median

A1ft: ft their s_x or s_x . NB using s gives awrt14.3

(d) M1: For a suitable comparison following through their value for the median of t.

A1: A correct conclusion in context following through their value for the median of t.

8MA0/02: AS Paper 2 Part A Statistics Mark scheme

Question	Sche	eme	Marks	AOs
2(a)	P(5£ X < 12) = P(X£ 11) - P(X£ 4)		M1	1.1b
	= 0.8939 - 0.0495	= awrt <u>0.844</u>	A1	1.1b
			(2)	
(b)	H_0 : $p = 0.25$ H_1 : $p > 0.25$ (both	correct in terms of p or p	B1	2.5
	<i>Y</i> ∼B(40, 0.25)		M1	3.3
	Method 1	Method 2		
	$P(Y^3 16) = 1 - P(Y \pounds 15)$	$P(Y^3 17) = 0.0116$	M1	1.1b
	= 1- 0.9738	$P(Y^3 18) = 0.0047$		
	= 0.0262	CR Y ³ 18	A1	1.1b
	0.0262 > 0.01 16 < 18 or 16 is not in the critical region or 16 is not significant, accept H ₀ . There is no significant evidence that the proportion of people who bought organic eggs has increased		A1cso	2.2b
			(5)	
(c)	There is evidence that the proportion eggs has increased [since $0.05 > 0.05$]		B1ft	2.2b
			(1)	

(8 marks)

Notes:

(a)M 1: For dealing with P(5£ X < 12) they need to use the cumulative prob. Function on the calc.

A1: awrt 8.44 (from calculator).

(b) B1: Both hypotheses correct using p or p and 0.25

M1: Realising that the model B(40, 0.25) is to be used. This may be stated or used.

M1: Using or writing 1- $P(Y \pm 15)$ or 1- P(Y < 16)

a correct CR or $P(Y^3 \ 17) = 0.0116$ and $P(Y^3 \ 18) = 0.0047$

A1: awrt 0.0262 or CR Y^3 18 or Y > 17

A1cso: A fully correct solution with a correct conclusion in context to include the idea of proportion and increased plus referring to organic

(c) **B1ft:** For 0.0262 < 0.05 [ft their probability in part(b)] **or** a CR of 16^3 15 (allow 16 > 14) and a correct contextual conclusion.

Question	Scheme	Marks	AOs
3(a)	Pressure outliers are <1004.75 and >1018.75 Rainfall outliers are (<-3.05) and >82.95	M1	2.1
	p = 1019 and 1022 are outliers w = 102.0 is an outlier	Alcso	1.1b
		(2)	
(b)	e.g. was a negative correlation, now no (zero) correlation	B1	2.2b
		(1)	
(c)	e.g. there are a lot of zeros for rainfall in Perth and there are none in the sample. or e.g. these are the highest figures and you are unlikely to get these if the sample was random.	B1	2.4
		(1)	
(d)	On average, an increase of 1 hPa in daily mean pressure results in a decrease of 0.223 mm in daily rainfall.	B1	3.4
		(1)	
(e)	Unreliable, as the large data set does not cover December.	B1	2.4
		(1)	

(6 marks)

Notes:

(a) M1: At least one correct boundary point

A1: both upper boundary points and correct conclusions

- (b) **B1ft:** A suitable description of correlation before and after.
- (c) **B1:** For a comment that supports the idea that the sample is unlikely to be random.
- (d) **B1:** For a suitable description of the rate: rainfall per number of hPa with reference to figures
- (e) **B1:** For correct conclusion with a reason explaining why it would be extrapolation.

NB: B0 For out of range, extrapolation o.e. on their own without a reason.

Question	Scheme	Marks	AOs
4(a)	S and A since there is no intersection between A and S or the probability of S and A happening is zero	B1	1.2
		(1)	
(b)	(0.1+ p)' 0.25 = 0.1 [p = 0.3]	M1	3.1b
	q = 0.15 or 1 - q = 0.85	M1	1.1b
	r=1-"p"-"q"-0.25	M1dd	3.1b
	= 0.3	A1	1.1b
		(4)	
(c)	Independent since $0.25 \times "0.2" = 0.05$	B1	2.2a
		(1)	
(d)	The teacher's belief would appear not to be justified as <i>D</i> and <i>S</i> are independent	B1ft	2.4
		(1)	

(7 marks)

Notes:

(a) **B1:** For S and A and a sensible reason

(b) M1: For forming a correct equation in terms of p using the information given.

M1: Writing or using q = 0.15 or 1 - q = 0.85

M1dd: dependent on both previous M marks being awarded. For using their values for p and q to form a correct equation to enable them to find r

A1: cao

- (c) **B1:** Yes and a suitable reason to support their answer bringing together the two pieces of information to draw the correct conclusion
- (d) **B1:** A correct comment following their answer to part (c) with reference to the teachers belief.

8MA0/02: AS Paper 2 Part B Mechanics Mark scheme

Question	Scheme	Marks	AOs
1(a)	Use of $s = vt - \frac{1}{2}at^2$	M1	2.1
	$19.6 = 4v - \frac{1}{2} \cdot 9.8 \cdot 4^2$	A1	1.1b
	$v = 24.5 \text{ or } 25 \text{ (m s}^{-1})$	A1	1.1b
		(3)	
(b)	$0 = 14.7^2 - 2^9.8h$	M1	2.1
	h = 11.0 or 11 (m)	A1	1.1b
		(2)	
(c)	New value of speed would be lower.	B1	3.5a
		(1)	

(6 marks)

Notes:

(a)

M1: Complete method to give equation in v only (could involve 2 or more *suvat* equations and then elimination) with usual rules

A1: Correct equation

A1: Correct answer

(b)

M1: Complete method to find h

A1: 11.0 or 11 (m)

(c)

B1: New value of speed will be lower

Question	Scheme	Marks	AOs
2(a)	Shape 120	B1	1.1b
	V, 120	B1	1.1b
		(2)	
(b)	$\frac{1}{2}$ 120 $V = 1500$	M1	3.1b
	V = 25	A1	1.1b
		(2)	
(c)	Area of triangle = Distance travelled = $(\frac{1}{2} \cdot 120V) = 1500$	B1	2.4
	This does not depend on T so T can take any value where $0 < T < 120$	B1	2.4
		(2)	
(d)	Include a constant speed phase in the motion	B1	3.5c
		(1)	

(7 marks)

Notes:

(a)

B1: Triangle, starting at the origin with base on axis and apex between t = 0 and t = 120

B1: *V* and 120 correctly marked (allow a delineator)

(b)

M1: Identifying correct strategy to solve problem to give equation in V only

A1: V = 25

(c)

B1: Area of triangle only depends on base and height

B1: So *T* can take any value $0 \le T \le 120$

(d)

B1: e.g. Include a *smooth* change from acceleration phase to deceleration phase.

e.g. Have a variable acceleration and/or deceleration phase

Question	Scheme	Marks	AOs
3(a)(i)	Equation of motion for <i>P</i> with usual rules	M1	3.3
	$T-1.5=0.4\times2.5$	A1	1.1b
	T = 2.5 (N)	A1	1.1b
(ii)	Equation of motion for Q with usual rules	M1	3.3
	10M - T = 2.5M	A1	1.1b
	M = 0.33	A1	1.1b
		(6)	
(b)	$2 = \frac{1}{2} \times 2.5t^2$	M1	3.4
	t = 1.3 (s)	A1	1.1b
		(2)	
(c)	e.g. the mass of the rope	B1	3.5b
		(1)	

(9 marks)

Notes:

(a) (i)

M1: Resolve horizontally for P

A1: Correct equation

A1: Correct answer. Ignore units

(a)(ii)

M1: Resolve vertically for Q

A1: Correct equation

A1: Correct answer

(b)

M1: Use $s = ut + \frac{1}{2}at^2$

A1: 1.3. Ignore units

(c)

B1: e.g. the pulley may not be smooth, air resistance

Question	Scheme	Marks	AOs
4(a)	$s = \oint_0^1 16 - 3t^2 dt$	M1	1.1a
	$= \left[16t - t^3\right]_0^1$	A1	1.1b
	=15 (m)	A1	1.1b
		(3)	
(b)	$16 - 3t^2 = 0$	M1	3.1b
	$t = \sqrt{\frac{16}{3}} \text{ oe}$	A1	1.1b
		(2)	
(c)	$16t - t^3 = 0$	M1	3.1b
	$t(16 - t^2) = 0$	M1	1.1b
	t=4	A1	1.1b
		(3)	

(8 marks)

Notes:

(a)

M1: Attempt to integrate, one power going up

A1: Correct integral and limits or indefinite integral with C = 0 and t = 1.

A1: 15 (m)

(b)

M1: Identifying correct strategy to solve problem of finding direction change by equating v to 0 and solving for t

A1: correct answer – any surd or decimal equivalent to at least 2 sf

(c)

M1: Identifying correct strategy to solve problem by using use s = 0 and equating their integral to 0

M1: Attempt to solve

A1: t = 4