



Maths Questions By Topic:

Statistical Sampling

A-Level Edexcel

 0207 060 4494

 www.expert-tuition.co.uk

 online.expert-tuition.co.uk

 enquiries@expert-tuition.co.uk

 The Foundry, 77 Fulham Palace Road, W6 8JA

Table Of Contents

New Spec

Paper 2 (AS) Page 1

Paper 3 (A2) Page 7

Old Spec

Statistics 2 Page 9

1. Helen is studying one of the qualitative variables from the large data set for Heathrow from 2015.

She started with the data from 3rd May and then took every 10th reading.

There were only 3 different outcomes with the following frequencies

Outcome	<i>A</i>	<i>B</i>	<i>C</i>
Frequency	16	2	1

(a) State the sampling technique Helen used. (1)

(b) From your knowledge of the large data set

- (i) suggest which variable was being studied,
- (ii) state the name of outcome *A*.

(2)

George is also studying the same variable from the large data set for Heathrow from 2015. He started with the data from 5th May and then took every 10th reading and obtained the following

Outcome	<i>A</i>	<i>B</i>	<i>C</i>
Frequency	16	1	1

Helen and George decided they should examine all of the data for this variable for Heathrow from 2015 and obtained the following

Outcome	<i>A</i>	<i>B</i>	<i>C</i>
Frequency	155	26	3

(c) State what inference Helen and George could reliably make from their original samples about the outcomes of this variable at Heathrow, for the period covered by the large data set in 2015. (1)

2. A lake contains three different types of carp.

There are an estimated 450 mirror carp, 300 leather carp and 850 common carp.

Tim wishes to investigate the health of the fish in the lake.

He decides to take a sample of 160 fish.

(a) Give a reason why stratified random sampling cannot be used.

(1)

(b) Explain how a sample of size 160 could be taken to ensure that the estimated populations of each type of carp are fairly represented.

You should state the name of the sampling method used.

(2)

(Total for Question 2 is 3 marks)

4. Sara is investigating the variation in daily maximum gust, t kn, for Camborne in June and July 1987.

She used the large data set to select a sample of size 20 from the June and July data for 1987. Sara selected the first value using a random number from 1 to 4 and then selected every third value after that.

(a) State the sampling technique Sara used. (1)

(b) From your knowledge of the large data set explain why this process may not generate a sample of size 20. (1)

(Total for Question 4 is 2 marks)

5. Sara was studying the relationship between rainfall, r mm, and humidity, $h\%$, in the UK. She takes a random sample of 11 days from May 1987 for Leuchars from the large data set.

She obtained the following results.

h	93	86	95	97	86	94	97	97	87	97	86
r	1.1	0.3	3.7	20.6	0	0	2.4	1.1	0.1	0.9	0.1

Sara examined the rainfall figures and found

$$Q_1 = 0.1 \quad Q_2 = 0.9 \quad Q_3 = 2.4$$

A value that is more than 1.5 times the interquartile range (IQR) above Q_3 is called an outlier.

- (i) Comment on the suitability of Sara's sampling method for this study. (2)
- (ii) Suggest how Sara could make better use of the large data set for her study.

(Total for Question 5 is 2 marks)

