

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

---

Forename(s)

---

Candidate signature

---

I declare this is my own work.

# A-level CHEMISTRY

## Paper 1 Inorganic and Physical Chemistry

Tuesday 2 June 2020

Afternoon

Time allowed: 2 hours

### Materials

For this paper you must have:

- the Periodic Table/Data Booklet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

### Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 105.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
<b>TOTAL</b>	



Answer **all** questions in the spaces provided.


0 1

This question is about enthalpy changes.

0 1 . 1

**Figure 1** shows a Born–Haber cycle for the formation of strontium chloride,  $\text{SrCl}_2$

**Figure 1**



**Table 1** shows some thermodynamic data.

**Table 1**

	Enthalpy change / $\text{kJ mol}^{-1}$
First ionisation energy of strontium	+548
Second ionisation energy of strontium	+1060
Enthalpy of atomisation of chlorine	+121
Enthalpy of atomisation of strontium	+164
Enthalpy of formation of strontium chloride	−828
Enthalpy of lattice formation of strontium chloride	−2112



Use the data in **Table 1** to calculate a value for the electron affinity of chlorine.

**[3 marks]**

Electron affinity \_\_\_\_\_  $\text{kJ mol}^{-1}$

**0 1 . 2** Draw a line from **each** substance to the enthalpy of lattice formation of that substance.  
**[1 mark]**

Substance	Enthalpy of lattice formation / $\text{kJ mol}^{-1}$
<input type="text" value="MgCl&lt;sub&gt;2&lt;/sub&gt;"/>	<input type="text" value="-2018"/>
<input type="text" value="MgO"/>	<input type="text" value="-2493"/>
<input type="text" value="BaCl&lt;sub&gt;2&lt;/sub&gt;"/>	<input type="text" value="-3889"/>

Question 1 continues on the next page

Turn over ►



**Table 2** shows the theoretical lattice enthalpy, based on a perfect ionic model, and an experimental value for the enthalpy of lattice formation of silver chloride.

**Table 2**

	Theoretical	Experimental
Enthalpy of lattice formation / $\text{kJ mol}^{-1}$	-770	-905

**0 1 . 3** State why there is a difference between the theoretical and experimental values. **[1 mark]**

---



---

**0 1 . 4** **Table 3** shows enthalpy of hydration values for ions of some Group 1 elements.

**Table 3**

	$\text{Li}^+(\text{g})$	$\text{Na}^+(\text{g})$	$\text{K}^+(\text{g})$
Enthalpy of hydration / $\text{kJ mol}^{-1}$	-519	-406	-322

Explain why the enthalpy of hydration becomes less exothermic from  $\text{Li}^+$  to  $\text{K}^+$  **[2 marks]**

---



---



---



---



0 1 . 5 Calcium bromide dissolves in water.

**Table 4** shows some enthalpy data.

**Table 4**

	Enthalpy change / $\text{kJ mol}^{-1}$
Enthalpy of solution of calcium bromide	-110
Enthalpy of lattice formation of calcium bromide	-2176
Enthalpy of hydration of calcium ions	-1650

Use the data in **Table 4** to calculate the enthalpy of hydration, in  $\text{kJ mol}^{-1}$ , of bromide ions.

**[3 marks]**

Enthalpy of hydration of bromide ions \_\_\_\_\_  $\text{kJ mol}^{-1}$

10

**Turn over for the next question**

**Turn over ►**



**0 2**

This question is about the isotopes of chromium.

**0 2 . 1**

Give the meaning of the term relative atomic mass.

**[2 marks]**

---

---

---

---

**0 2 . 2**A sample of chromium containing the isotopes  $^{50}\text{Cr}$ ,  $^{52}\text{Cr}$  and  $^{53}\text{Cr}$  has a relative atomic mass of 52.1The sample contains 86.1% of the  $^{52}\text{Cr}$  isotope.

Calculate the percentage abundance of each of the other two isotopes.

**[4 marks]**Abundance of  $^{50}\text{Cr}$  \_\_\_\_\_ %      Abundance of  $^{53}\text{Cr}$  \_\_\_\_\_ %

0 2 . 3

State, in terms of the numbers of fundamental particles, **one** similarity and **one** difference between atoms of  $^{50}\text{Cr}$  and  $^{53}\text{Cr}$

**[2 marks]**

Similarity \_\_\_\_\_

\_\_\_\_\_

Difference \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

The sample of chromium is analysed in a time of flight (TOF) mass spectrometer.

0 2 . 4

Give **two** reasons why it is necessary to ionise the isotopes of chromium before they can be analysed in a TOF mass spectrometer.

**[2 marks]**

1 \_\_\_\_\_

\_\_\_\_\_

2 \_\_\_\_\_

\_\_\_\_\_

**Question 2 continues on the next page**

**Turn over ►**

0 2 . 5

A  $^{53}\text{Cr}^+$  ion travels along a flight tube of length 1.25 m  
The ion has a constant kinetic energy ( $KE$ ) of  $1.102 \times 10^{-13}$  J

$$KE = \frac{mv^2}{2}$$

$m$  = mass of the ion / kg

$v$  = speed of ion /  $\text{m s}^{-1}$

Calculate the time, in s, for the  $^{53}\text{Cr}^+$  ion to travel down the flight tube to reach the detector.

The Avogadro constant,  $L = 6.022 \times 10^{23} \text{ mol}^{-1}$

**[5 marks]**

Time \_\_\_\_\_ s

15





**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**




0 3

This question is about Period 3 elements.

**Figure 2** shows the **second** ionisation energies of some elements in Period 3.

**Figure 2**



0 3 . 1

Draw a cross (x) on **Figure 2** to show the **second** ionisation energy of silicon.

[1 mark]

0 3 . 2

Identify the element in Period 3, from sodium to argon, that has the highest **second** ionisation energy.

Give an equation, including state symbols, to show the process that occurs when the **second** ionisation energy of this element is measured.

If you were unable to identify the element you may use the symbol **Q** in your equation.

[2 marks]

Element \_\_\_\_\_

Equation

\_\_\_\_\_

0 3 . 3

Explain why the atomic radius decreases across Period 3, from sodium to chlorine.

[2 marks]

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_



0 3 . 4

Identify the element in Period 3, from sodium to chlorine, that has the highest electronegativity.

**[1 mark]**

0 3 . 5

Phosphorus burns in air to form phosphorus(V) oxide.  
Give an equation for this reaction.

**[1 mark]**

---

**7****Turn over for the next question****Turn over ►**

**0 4** Propanoic acid ( $\text{C}_2\text{H}_5\text{COOH}$ ) is a weak acid.

The acid dissociation constant ( $K_a$ ) for propanoic acid is  $1.35 \times 10^{-5} \text{ mol dm}^{-3}$  at  $25^\circ\text{C}$

**0 4 . 1** State the meaning of the term weak acid.

[1 mark]

---

---

**0 4 . 2** Give an expression for the acid dissociation constant for propanoic acid.

[1 mark]

$K_a$

**0 4 . 3** A student dilutes  $25.0 \text{ cm}^3$  of  $0.500 \text{ mol dm}^{-3}$  propanoic acid by adding water until the total volume is  $100.0 \text{ cm}^3$

Calculate the pH of this diluted solution of propanoic acid.

Give your answer to 2 decimal places.

[4 marks]

pH \_\_\_\_\_



0 4 . 4

A buffer solution with a pH of 4.50 is made by dissolving  $x$  g of sodium propanoate ( $\text{C}_2\text{H}_5\text{COONa}$ ) in a solution of propanoic acid. The final volume of buffer solution is  $500 \text{ cm}^3$  and the final concentration of the propanoic acid is  $0.250 \text{ mol dm}^{-3}$

Calculate  $x$  in g

For propanoic acid,  $K_a = 1.35 \times 10^{-5} \text{ mol dm}^{-3}$

[6 marks]

$x$  \_\_\_\_\_ g


12

Turn over ►



0 5

Some reactions of the  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}(\text{aq})$  ion are shown.



0 5 . 1

Give the formula of the white precipitate **B**.

State **one** other observation when  $\text{Na}_2\text{CO}_3(\text{aq})$  is added to a solution containing  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}(\text{aq})$  ions.

Give an equation for this reaction.

[3 marks]

Formula of **B** \_\_\_\_\_

Observation \_\_\_\_\_

Equation

\_\_\_\_\_

0 5 . 2

Give the formula of the complex ion **C**.

State **one** condition needed for the formation of **C** from  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}(\text{aq})$  and  $\text{NaOH}(\text{aq})$ .

Give an equation for this reaction.

[3 marks]

Formula of **C** \_\_\_\_\_

Condition \_\_\_\_\_

Equation

\_\_\_\_\_



**0 5** . **3** Deduce the formula of the complex ion **A**.

**[1 mark]**

---

**0 5** . **4** Explain, with the use of an equation, why a solution containing  $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$  has a pH < 7

**[3 marks]**

Equation

---

Explanation

---

---

---

---

10


**Turn over for the next question**

**Turn over ►**




0 6

Methanol can be manufactured in a reversible reaction as shown.



**Figure 3** shows how the partial pressures change with time at a constant temperature.

**Figure 3**



0 6

1

Draw a cross (x) on the appropriate axis of **Figure 3** when the mixture reaches equilibrium.

[1 mark]

0 6

2

A 0.230 mol sample of carbon monoxide is mixed with hydrogen in a 1:2 mol ratio and allowed to reach equilibrium in a sealed flask at temperature  $T$ . At equilibrium the mixture contains 0.120 mol of carbon monoxide. The total pressure of this mixture is  $1.04 \times 10^4$  kPa

Calculate the partial pressure, in kPa, of hydrogen in the equilibrium mixture.

[4 marks]

Partial pressure of hydrogen \_\_\_\_\_ kPa





**0 6 . 3** Give an expression for the equilibrium constant ( $K_p$ ) for this reaction.

State the units.

**[2 marks]**

$K_p$

Units \_\_\_\_\_

**0 6 . 4** Some more carbon monoxide is added to the mixture in Question **06.2**. The new mixture is allowed to reach equilibrium at temperature  $T$ .

State the effect, if any, on the partial pressure of methanol and on the value of  $K_p$

**[2 marks]**

Effect on partial pressure of methanol \_\_\_\_\_

Effect on value of  $K_p$  \_\_\_\_\_

**0 6 . 5** State the effect, if any, of the addition of a catalyst on the value of  $K_p$  for this equilibrium.  
Explain your answer.

**[2 marks]**

Effect on value of  $K_p$  \_\_\_\_\_

Explanation \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

11

**Turn over for the next question**

**Turn over ►**







**0 8**

A student does an experiment to determine the percentage by mass of sodium chlorate(I), NaClO, in a sample of bleach solution.



Method:

- Dilute a 10.0 cm<sup>3</sup> sample of bleach solution to 100 cm<sup>3</sup> with distilled water.
- Transfer 25.0 cm<sup>3</sup> of the diluted bleach solution to a conical flask and acidify using sulfuric acid.
- Add excess potassium iodide to the conical flask to form a brown solution containing I<sub>2</sub>(aq).
- Add 0.100 mol dm<sup>-3</sup> sodium thiosulfate solution (Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>) to the conical flask from a burette until the brown solution containing I<sub>2</sub>(aq) becomes a colourless solution containing I<sup>-</sup>(aq).

The student uses 33.50 cm<sup>3</sup> of sodium thiosulfate solution.

The density of the original bleach solution is 1.20 g cm<sup>-3</sup>

The equations for the reactions in this experiment are

**0 8 . 1**

Use all the information given to calculate the percentage by mass of NaClO in the original bleach solution.

Give your answer to 3 significant figures.

**[7 marks]**

Percentage by mass \_\_\_\_\_



0 8 . 2

The total uncertainty from two readings and an end point error in using a burette is  $\pm 0.15 \text{ cm}^3$

What is the total percentage uncertainty in using the burette in this experiment?

**[1 mark]**

Tick (✓) **one** box.

0.45%

0.90%

1.34%

---

8

**Turn over for the next question**

**Turn over ►**

0 9

This question is about sodium halides.

0 9 . 1

State what is observed when silver nitrate solution is added to sodium fluoride solution.

**[1 mark]**

---

0 9 . 2

State **one** observation when solid sodium chloride reacts with concentrated sulfuric acid.

Give an equation for the reaction.

State the role of the chloride ions in the reaction.

**[3 marks]**Observation 

---

Equation 

---

Role 

---

0 9 . 3

Give an equation for the redox reaction between solid sodium bromide and concentrated sulfuric acid.

Explain, using oxidation states, why this is a redox reaction.

**[3 marks]**Equation 

---

Explanation 

---

---

---

---

0 9 . 4

State what is observed when aqueous chlorine is added to sodium bromide solution.

Give an ionic equation for the reaction.

**[2 marks]**Observation 

---

Ionic equation 

---


---

---

  
9

1 0

Methanol is formed when carbon dioxide and hydrogen react.



**Table 5** contains enthalpy of formation and entropy data for these substances.

**Table 5**

	<b>CO<sub>2</sub>(g)</b>	<b>H<sub>2</sub>(g)</b>	<b>CH<sub>3</sub>OH(g)</b>	<b>H<sub>2</sub>O(g)</b>
$\Delta_f H / \text{kJ mol}^{-1}$	-394	0	-201	-242
$S / \text{J K}^{-1} \text{mol}^{-1}$	214	131	238	189

1 0 . 1

Use the equation and the data in **Table 5** to calculate the Gibbs free-energy change ( $\Delta G$ ), in  $\text{kJ mol}^{-1}$ , for this reaction at 890 K

**[6 marks]**

$\Delta G$  \_\_\_\_\_  $\text{kJ mol}^{-1}$


**Turn over ►**



**Figure 4** shows how the Gibbs free-energy change varies with temperature in a different gas phase reaction.

The straight line graph for this gas phase reaction has been extrapolated to zero Kelvin.

**Figure 4**





1 0 . 2

Use the values of the intercept and gradient from the graph in **Figure 4** to calculate the enthalpy change ( $\Delta H$ ), in  $\text{kJ mol}^{-1}$ , and the entropy change ( $\Delta S$ ), in  $\text{J K}^{-1} \text{mol}^{-1}$ , for this reaction.

**[4 marks]** $\Delta H$  \_\_\_\_\_  $\text{kJ mol}^{-1}$  $\Delta S$  \_\_\_\_\_  $\text{J K}^{-1} \text{mol}^{-1}$ 

1 0 . 3

State what **Figure 4** shows about the feasibility of the reaction.

**[1 mark]**

---

---

---

11

**Turn over ►**

1	1
---	---

This question is about a glucose–oxygen fuel cell.

When the cell operates, the glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ) molecules react with water at the negative electrode to form carbon dioxide and hydrogen ions.

Oxygen gas reacts with hydrogen ions to form water at the positive electrode.

1	1	.	1
---	---	---	---

Deduce the half-equation for the reaction at the negative electrode.

[1 mark]

---

1	1	.	2
---	---	---	---

Deduce the half-equation for the reaction at the positive electrode.

[1 mark]

---

1	1	.	3
---	---	---	---

Give the equation for the overall reaction that occurs in the Glucose–oxygen fuel cell.

[1 mark]

---

1	1	.	4
---	---	---	---

The negative electrode is made of carbon and the positive electrode is made of platinum.

Give the conventional representation for the glucose–oxygen fuel cell.

[2 marks]

---

---

1	1	.	5
---	---	---	---

State what must be done to maintain the EMF of this fuel cell when in use.

[1 mark]

---

6
---

**END OF QUESTIONS**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**











**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.aqa.org.uk](http://www.aqa.org.uk).

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2020 AQA and its licensors. All rights reserved.



3 2



2 0 6 A 7 4 0 5 / 1

IB/M/Jun20/7405/1