



# Mark Scheme (Results)

Summer 2019

Pearson GCE Advanced Subsidiary Level  
In Chemistry (8CH0) Paper 01 Core Inorganic  
and Physical Chemistry

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk) for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson. Their contact details can be found on this link: [www.edexcel.com/teachingservices](http://www.edexcel.com/teachingservices).

You can also use our online Ask the Expert service at [www.edexcel.com/ask](http://www.edexcel.com/ask). You will need an Edexcel username and password to access this service.

### **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2019

Publications Code 8CH0\_01\_1906\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
  - i) ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
  - ii) select and use a form and style of writing appropriate to purpose and to complex subject matter
  - iii) organise information clearly and coherently, using specialist vocabulary when appropriate

## Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

( ) means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

### Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Question Number	Answer	Mark
1	<p><b>The only correct answer is D</b> (<math>9.03 \times 10^{24}</math>)</p> <p><i>A is not correct because this is the answer for 1 mol of aluminium oxide as molecules</i></p> <p><i>B is not correct because this is the answer for 3 mol of aluminium oxide as molecules</i></p> <p><i>C is not correct because this is the answer for the ions in 1 mol of aluminium oxide</i></p>	(1)

(Total for Question 1 = 1 mark)

Question Number	Answer	Mark
2	<p><b>The only correct answer is D</b> (<math>1s^2, 2s^2, 2p^6, 3s^2, 3p^6</math>)</p> <p><i>A is not correct because <math>1s^2, 2s^2, 2p^6, 3s^2, 3p^2</math> is for an <math>S^{2+}</math> ion</i></p> <p><i>B is not correct because <math>1s^2, 2s^2, 2p^6, 3p^4</math> is for an <math>S^{2+}</math> ion with electrons removed from the 3s subshell</i></p> <p><i>C is not correct because <math>1s^2, 2s^2, 2p^6, 3s^2, 3p^4</math> is for the sulfur atom</i></p>	(1)

(Total for Question 2 = 1 mark)

Question Number	Acceptable Answer	Additional Guidance	Mark
3(a)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• same number of protons (1)</li> <li>• different numbers of neutrons (1)</li> </ul>	<p>Ignore number of electrons</p> <p>Ignore references to atom(s) / 'elements' in the answer</p>	(2)


Question Number	Acceptable Answer	Additional Guidance	Mark
3(b)	<ul style="list-style-type: none"> <li>• correct subtraction to calculate relative abundance of unknown isotope (1)</li> <li>• calculation of mass number of unknown isotope of gallium with suitable working (1)</li> </ul>	<p><u>Example of calculation</u>  <math>100 - 63.25 = 36.75</math>  <b>OR</b>  <math>1.00 - 0.6325 = 0.3675</math></p> <p><math>\frac{(69 \times 63.25) + (M \times 36.75)}{100} = 69.735</math></p> <p>OR</p> <p><math>43.64(25) + \frac{36.75 M}{100} = 69.735</math></p> <p><math>M = 71</math></p> <p>Ignore any units given with final answer.  Correct answer with no working gets M2 only  Allow TE from M1</p>	(2)

(Total for Question 3 = 4 marks)

Question Number	Answer	Mark
4(a)	<p><b>The only correct answer is A</b> (carbonates – increases, nitrates – increases)</p> <p><i>B is not correct because thermal stability of Group 2 nitrates does not decrease down the group</i></p> <p><i>C is not correct because thermal stability of Group 2 carbonates does not decrease down the group</i></p> <p><i>D is not correct because thermal stabilities of Group 2 carbonates and nitrates do not decrease down the group</i></p>	<b>(1)</b>





	<ul style="list-style-type: none"> <li>• safety precaution: fume cupboard/hood (1)</li> </ul>	<p>Example diagram:</p>  <p>Ignore well ventilated room / face mask / goggles / gloves / lab coat This is the only acceptable safety precaution.</p>	
--	---	---	--

Question Number	Answer	Mark
4(c)	<p><b>The only correct answer is B</b> (hydroxides – increases, sulfates – decreases)</p> <p><i>A is not correct because sulfate solubility does not increase down the group</i></p> <p><i>C is not correct because hydroxide solubility does not decrease down the group and sulfate solubility does not increase down the group</i></p> <p><i>D is not correct because hydroxide solubility does not decrease down the group</i></p>	(1)

(Total for Question 4 = 6 marks)

Question Number	Answer	Mark
5(a)	<p><b>The only correct answer is C (36.7%)</b></p> <p><i>A is not correct because 21.3% is calculated using the atomic number of iron</i></p> <p><i>B is not correct because 35.1% is calculated using all atomic numbers</i></p> <p><i>D is not correct because 53.8% is calculated using the atomic numbers of sulfur and oxygen</i></p>	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
5(b)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(add hydrochloric acid / nitric acid then) add barium chloride / barium nitrate (solution)      <b>(1)</b></li> <li>white precipitate / white solid      <b>(1)</b></li> </ul>	<p>Ignore omission of acid Do not award just Barium ions / Ba<sup>2+</sup></p> <p>M2 is dependent on M1, with the exception of just Ba<sup>2+</sup> given as reagent</p>	(2)

Question Number	Answer	Mark
5(c)	<p><b>The only correct answer is A (392.0)</b></p> <p><i>B is not correct because 312.0 is calculated from only 6 multiples of the H<sub>2</sub> of the 6-water</i></p> <p><i>C is not correct because 302.0 is calculated by not multiplying the water by 6</i></p> <p><i>D is not correct because 284.0 is calculated by ignoring the 6-water completely</i></p>	(1)

**(Total for Question 5 = 4 marks)**

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li data-bbox="401 402 1108 435">• any mention of platinum/nichrome wire <b>(1)</b></li> <li data-bbox="401 630 1171 695">• dip the wire into (clean/fresh concentrated) hydrochloric acid <b>(1)</b></li> <li data-bbox="401 857 1108 922">• dip the (wet) wire into the solid and place in a (non-luminous/roaring/blue Bunsen) flame <b>(1)</b></li> </ul>	<p>Allow NiCr for nichrome  Allow silica/magnesia for platinum or nichrome  Allow loop / rod for wire  Ignore inoculating / flame-test (wire)</p> <p>Allow any mention of HCl(aq) e.g. cleaning or mixing solid and acid or making a paste  Allow HCl for HCl(aq)</p> <p>Ignore dilute</p> <p>Allow on / over / under / near / show / above for 'in'</p>	<b>(3)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(ii)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• mention of energy or heat or heating (from the flame) <b>(1)</b></li> <li>• electrons promoted to higher energy levels <b>(1)</b></li> <li>• electrons drop down / return (to lower energy levels / ground state) <b>(1)</b></li> <li>• light (in the visible region) is emitted / released / given out <b>(1)</b></li> </ul>	<p>Do not award M1 for “burning”</p> <p>Allow just ‘electrons excited’ for M2</p> <p>Allow electromagnetic / e.m. radiation / photons instead of light</p>	<b>(4)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(iii)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>• no emission of light /energy in the visible region (of the spectrum)</li> </ul>	<p>Do not award any mention of (bright) white light emission</p> <p>Allow electromagnetic / e.m. radiation / photons / colour instead of light / energy</p> <p>Allow the light emitted is in the UV or IR</p> <p>Allow any references to frequency or wavelength being too high or too low</p>	<b>(1)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
6(a)(iv)	An answer that makes reference to the following point: <ul style="list-style-type: none"> <li>(There are) other ions that do not produce a flame colour</li> </ul>	Allow a specific ion that does not have a flame colour e.g. 'beryllium' / $\text{Be}^{2+}$ Allow other "elements" do not produce a flame colour Do not award if any references to "burning"	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
6(b)(i)	<ul style="list-style-type: none"> <li><math>\text{AgNO}_3</math></li> </ul>	Ignore 'silver nitrate' Ignore state symbols for $\text{AgNO}_3$	(1)

Question Number	Answer	Mark
6(b)(ii)	<p><b>The only correct answer is B</b> (insoluble in dilute ammonia solution, soluble in concentrated ammonia solution)</p> <p><i>A is not correct because silver bromide is soluble in concentrated ammonia solution</i></p> <p><i>C is not correct because silver bromide is insoluble in dilute ammonia solution and soluble in concentrated ammonia solution</i></p> <p><i>D is not correct because silver bromide is insoluble in dilute ammonia solution</i></p>	(1)

(Total for Question 6 = 11 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(a)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>balanced equation with correct species <b>(1)</b></li> <li>correct states all correct <b>(1)</b></li> </ul>	<p><u>Example of equation:</u></p> $\text{Mg(s)} + 2\text{HCl(aq)} \rightarrow \text{MgCl}_2\text{(aq)} + \text{H}_2\text{(g)}$ <p>or</p> $\text{Mg(s)} + 2\text{H}^+\text{(aq)} \rightarrow \text{Mg}^{2+}\text{(aq)} + \text{H}_2\text{(g)}$ <p>Do not award M2 for incorrect formulae e.g. MgCl (for MgCl<sub>2</sub>), or H (for H<sub>2</sub>)</p> <p>Allow M2 for unbalanced equation if all species correct</p>	<b>(2)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
7(b)(i)	<p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>calculation of uncertainty</li> </ul>	<p><u>Example of calculation:</u></p> $\frac{(\pm)0.5 \times 100}{10.0} = (\pm)5/5.0/5.00(\%)$	<b>(1)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
7(b)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• calculation of moles of Mg <span style="float: right;">(1)</span></li> <li>• calculation of moles of HCl <span style="float: right;">(1)</span></li> <li>• evidence to support Mg in excess <span style="float: right;">(1)</span></li> </ul>	<p><u>Example of calculation:</u></p> <p><u>0.12</u>  <math>24.3 = 4.9383 \times 10^{-3} / 0.0049383</math> (mol)</p> <p>Allow <math>A_r</math> for Mg = 24</p> <p><u>10 x 0.20</u>  <math>\frac{1000}{1000} = 2.0 \times 10^{-3} / 0.002</math> (mol)</p> <p><math>4.9383 \times 10^{-3}</math> mol of Mg requires  <math>9.8765 \times 10^{-3}</math> mol of HCl          (and <math>0.002 &lt; 9.8 \times 10^{-3}</math>) so Mg in excess          or          0.002 mol HCl requires 0.001 mol Mg          (and <math>0.0049 &gt; 0.001</math>) so Mg in excess          Ignore SF</p> <p>Do not award <b>M3</b> for <math>0.0049 &gt; 2 \times 0.002</math>          OR  <math>0.0049 &gt; 0.004</math> to show that Mg is in excess</p> <p>Do not award M3 if HCl stated to be in excess</p>	(3)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(b)(iii)	calculation of moles of gas	<u>Example of calculation</u> $0.002 \div 2 = 0.001$ or $1 \times 10^{-3}$ Allow TE from (a) and (b)(ii)	<b>(1)</b>



Question Number	Acceptable Answer	Additional Guidance	Mark
7(b)(iv)	<ul style="list-style-type: none"> <li data-bbox="352 402 968 435">• Rearrangement of ideal gas equation (1)</li> <li data-bbox="352 589 968 621">• conversion of °C to K (1)</li> <li data-bbox="352 695 968 727">• calculation of volume in m<sup>3</sup> (1)</li> <li data-bbox="352 963 968 995">• calculation of volume in cm<sup>3</sup> (1)</li> </ul>	<p data-bbox="1045 329 1346 362"><u>Example of calculation:</u></p> <p data-bbox="1045 402 1398 475"><math>pV = nRT</math> rearrange <math>V = \frac{nRT}{p}</math></p> <p data-bbox="1045 475 1776 548">Allow M1 if equation rearrangement not explicitly shown but used correctly in M3</p> <p data-bbox="1045 589 1545 662">(273 + 23) = 296 Allow M2 if (273 + 23) used in equation</p> <p data-bbox="1045 703 1461 776"><math>V = \frac{1.0 \times 10^{-3} \times 8.31 \times (273 + 23)}{98\,000}</math></p> <p data-bbox="1045 816 1283 849">= 2.51 × 10<sup>-5</sup> (m<sup>3</sup>)</p> <p data-bbox="1045 889 1318 922">= 25 allow 25.1 (cm<sup>3</sup>)</p> <p data-bbox="1045 963 1577 1036">Allow TE from (b)(iii) and TE at each stage Allow <b>2 or 3</b> SF for final answer</p> <p data-bbox="1045 1076 1356 1109"><b>ECF values from (b)(iii)</b></p> <p data-bbox="1045 1109 1434 1141">For 0.002 mol H<sub>2</sub>, V = 50.2 cm<sup>3</sup></p> <p data-bbox="1045 1141 1461 1174">For 0.00494 mol H<sub>2</sub>, V = 124 cm<sup>3</sup></p> <p data-bbox="1045 1174 1461 1206">For 0.00894 mol H<sub>2</sub>, V = 224 cm<sup>3</sup></p> <p data-bbox="1045 1206 1434 1239">For 0.004 mol H<sub>2</sub>, V = 100 cm<sup>3</sup></p>	

Question Number	Acceptable Answer	Additional Guidance	Mark
7(c)(i)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>gas lost before the bung replaced (1)</li> <li>the magnesium was coated with oxide (so water was formed instead of hydrogen) (1)</li> </ul>	<p>Ignore 'generic' gas leakages from apparatus Do not award gas may dissolve (in water or acid)</p> <p>Ignore 'generic' references to impurity Ignore references to incomplete reaction</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
7(c)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>arrange equipment so that the Mg ribbon drops into the acid <b>after</b> the delivery tube was replaced (1)</li> <li>clean the magnesium ribbon (1)</li> </ul>	<p>Ignore replace the bung more quickly Allow any workable method</p>	(2)

(Total for Question 7 = 15 marks)

Question Number	Acceptable Answer	Additional Guidance	Mark
8(a)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li data-bbox="363 402 1144 440">• comparison of ionic charges <span style="float: right;">(1)</span></li> <li data-bbox="363 630 1144 667">• comparison of ionic radii <span style="float: right;">(1)</span></li> <li data-bbox="363 813 1144 850">• comparison of <b>energy</b> required <span style="float: right;">(1)</span></li> </ul>	<p><u>Examples of explanations</u></p> <p>MgO has doubly charged ions <b>and</b> KBr has singly charged ions Allow reference to just one ion in each compound</p> <p>Mg<sup>2+</sup> smaller than K<sup>+</sup> and/or O<sup>2-</sup> smaller than Br<sup>-</sup> Ignore references to <b>atomic</b> radii</p> <p>More <b>energy</b> needed to overcome the electrostatic attractions/bonds (between ions) in MgO (than in KBr) Ignore references to 'electronegativity' / (ion) polarisation Award <b>(0)</b> overall if any mention of any of the following: London Forces <b>Molecules</b> / intermolecular forces Hydrogen bonding Covalent bonding</p>	<b>(3)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
8(b)	<p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• (solid potassium bromide does not conduct because) the <b>ions</b> are in fixed positions / <b>ions</b> are not free to move (1)</li> <li>• it does conduct in solution because the <b>ions</b> are free to move (and carry charge) (1)</li> </ul>	<p>Do not award any marks if reference to movement of <b>electrons</b> or <b>free electrons</b> when conduction occurs</p> <p>Do not award any marks if 'molecules'/ London forces mentioned</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark												
*8(c)	<p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="365 586 1136 894"> <thead> <tr> <th data-bbox="365 586 751 699">Number of indicative marking points seen in answer</th> <th data-bbox="751 586 1136 699">Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td data-bbox="365 699 751 740">6</td> <td data-bbox="751 699 1136 740">4</td> </tr> <tr> <td data-bbox="365 740 751 781">5-4</td> <td data-bbox="751 740 1136 781">3</td> </tr> <tr> <td data-bbox="365 781 751 821">3-2</td> <td data-bbox="751 781 1136 821">2</td> </tr> <tr> <td data-bbox="365 821 751 862">1</td> <td data-bbox="751 821 1136 862">1</td> </tr> <tr> <td data-bbox="365 862 751 894">0</td> <td data-bbox="751 862 1136 894">0</td> </tr> </tbody> </table>	Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points	6	4	5-4	3	3-2	2	1	1	0	0	<p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning, scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p>	(6)
Number of indicative marking points seen in answer	Number of marks awarded for indicative marking points														
6	4														
5-4	3														
3-2	2														
1	1														
0	0														

The following table shows how the marks should be awarded for structure and lines of reasoning.

	Number of marks awarded for structure and sustained lines of reasoning
Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout.	2
Answer is partially structured with some linkages and lines of reasoning.	1
Answer has no linkages between points and is unstructured.	0

In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.


If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded do not deduct mark(s).

Comment: Look for the indicative marking points first, then consider the mark for the structure of the answer and sustained line of reasoning.


**Indicative content:**

- **IP1** graphene has a **single** layer / **single** sheet (of hexagons / rings)
- **IP2** graphene has delocalised electrons / electrons which are mobile
- **IP3** graphite has layers / sheets / planes **and** each carbon bonded to **three** others
- **IP4** graphite has delocalised electrons / electrons which are mobile (between the layers)
- **IP5** diamond has each carbon bonded to four other carbons / diamond has a tetrahedral arrangement (around each C atom) / tetrahedral structure
- **IP6** diamond's C atoms have all their outer / valence / fourth electrons involved in bonding  
**OR**  
diamond has no delocalised electrons / all electrons are localised

Allow annotated diagrams for all marking points



graphite



diamond

graphene



Allow 'free' for delocalised or mobile

Question Number	Acceptable Answer	Additional Guidance	Mark
8(d)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>iron atoms have greater mass than carbon atoms <b>(1)</b></li> <li>iron atoms pack closer than carbon atoms (in graphite) <b>(1)</b></li> </ul>	<p>Allow weigh more / higher <math>A_r</math> / greater molar mass</p> <p>Allow idea of more space (between the layers of atoms) in graphite</p> <p>Award (0) overall if mention of iron molecules or graphite molecules</p>	<b>(2)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
8(e)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>Lower value relates to (weak) London / van der Waals' forces (between the layers) <b>(1)</b></li> <li>Higher value refers to (strong) covalent (C-C) bonds (within each layer) <b>(1)</b></li> </ul>	<p>Allow 'pi-bonds (between layers)'</p> <p>Allow "(weak) intermolecular forces (between layers)"</p> <p>Allow (C-C) 'sigma bonds'</p>	<b>(2)</b>

**(Total for Question 8 = 15 marks)**



Question Number	Answer	Mark
9(a)(i)	<p><b>The only correct answer is C</b> (Disproportionation)</p> <p><i>A is not correct because oxidation and reduction are occurring</i></p> <p><i>B is not correct because oxidation and reduction are occurring</i></p> <p><i>D is not correct because two reactants are involved</i></p>	<b>(1)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
9(a)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(trigonal) pyramidal <b>(1)</b></li> <li>(predicted bond angle) 107° <b>(1)</b></li> <li>three groups / three pairs of bonding electrons and one lone pair <b>OR</b> lone pair – bond pair repulsion &gt; bond pair – bond pair repulsion <b>(1)</b></li> <li>(electron pairs / groups repel to positions of) minimum repulsion / maximum separation <b>(1)</b></li> </ul>	<p>For M1, this shape must be <b>named</b></p> <p>Allow answers in the range 106.5° to 107.5° (allow actual value 110°) Allow M2 on an annotated diagram</p> <p>Allow 'regions' for 'groups' or 'pairs'</p> <p>Allow statements such as "lone pair repulsion greater than bond pair repulsion"</p>	<b>(4)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
9(b)(i)	<ul style="list-style-type: none"> <li>• <math>4 \text{KClO}_3 \rightarrow 1 \text{KCl} + 3 \text{KClO}_4</math></li> </ul>	Allow just KCl with no number in front Allow multiples	(1)

Question Number	Acceptable Answer	Additional Guidance	Mark
9(b)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• use of <b>water only</b> as solvent (1)</li> <li>• add the mixture of solids / products to any one of the following :               <ul style="list-style-type: none"> <li>water only</li> <li><b>or</b></li> <li>ethanol only</li> <li><b>or</b></li> <li>a water plus ethanol mixture</li> <li><b>or</b></li> <li>water followed by ethanol</li> <li><b>or</b></li> <li>ethanol followed by water (1)</li> </ul> </li> <li>• <b>filter</b> off the undissolved potassium chlorate(VII) <math>\text{KClO}_4</math> (1)</li> </ul>	<p>Use of separating funnel or electrolysis scores (0) overall</p> <p>NB: If water <b>only</b> used as the solvent, both M1 and M2 are awarded</p> <p>Allow 'insoluble solid' / 'less soluble solid' / 'the precipitate' for <math>\text{KClO}_4</math>            Allow salt for solid</p>	(3)

Question Number	Acceptable Answer	Additional Guidance	Mark
9(c)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>conversion of water volume to cm<sup>3</sup> or g (1)</li> <li>calculation of the mass of chlorine in 2.5 x 10<sup>9</sup> g/cm<sup>3</sup> of water (1)</li> <li>conversion of the mass of chlorine to moles of chlorine gas (Cl<sub>2</sub>) by dividing by 71 (1)</li> </ul>	<p><u>Example of calculation:</u></p> $2\,500\text{ m}^3 = 2\,500 \times 10^6\text{ cm}^3 = 2.5 \times 10^9\text{ cm}^3$ <p><b>or</b></p> $= 2\,500 \times 10^6\text{ g} = 2.5 \times 10^9\text{ g}$ $\frac{2 \times 2.5 \times 10^9}{1 \times 10^6} = 5.0 \times 10^3\text{ g}$ $\frac{5.0 \times 10^3}{(35.5 \times 2)} = 70(.4)(\text{mol})$ <p>Allow alternative calculation methods Ignore SF for final answer Answer of 35.2 / 140.8 mol Cl<sub>2</sub> scores (2) marks</p>	(3)

Question Number	Acceptable Answer	Additional Guidance	Mark
9(d)	<p>An explanation that makes reference to the following points:</p> <p>Lower than 1 ppm</p> <ul style="list-style-type: none"> <li>• HClO will be low(er) <b>(1)</b></li> <li>• ineffective (as a disinfectant) <b>(1)</b></li> </ul> <p>Higher than 3 ppm</p> <ul style="list-style-type: none"> <li>• HCl will be high(er) <b>(1)</b></li> <li>• any relevant effect of increased HCl <b>(1)</b></li> </ul>	<p>Ignore reference to amount of Cl<sub>2</sub> being too low M2 dependent on correct M1</p> <p>M4 dependent on correct M3 Award effects including corrosive, alters or lowers pH NB: Do not award high(er) pH Award increases acidity / strongly acidic / toxicity Award any reasonable negative effect on swimmers e.g. irritation / irritant Ignore just 'harmful' / just 'dangerous' Ignore reference to amount of Cl<sub>2</sub> being too high and its effects</p>	<b>(4)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
9(e)(i)	<p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• diagram showing <b>both</b> Na<sup>+</sup> ion <b>and</b> ClO<sup>-</sup> ion surrounded by water molecules/solvated <b>(1)</b></li> <li>• correct orientation of the water molecules around <b>both</b> ions with a relevant dipole shown on at least one water molecule for each ion, (i.e. δ<sup>-</sup> on O for a water molecule next to Na<sup>+</sup> and a δ<sup>+</sup> on at least one H atom on a water molecule next to a ClO<sup>-</sup>) <b>(1)</b></li> </ul>	<p>Allow any number of water molecules (&gt;1) for <b>both</b> ions  For M1 to be awarded there must be more than one H<sub>2</sub>O molecule around each ion</p> <p>M2 can be awarded even if only one H<sub>2</sub>O molecule is shown next to each ion</p> <p>Allow one mark for one ion surrounded by correctly orientated water molecules.</p> <p>Written description only, covering the same two marking points scores one mark max</p> <p><b>Mention of hydrogen bonding or water drawn as "HO<sub>2</sub>" or NaClO shown as covalent scores (0) overall</b></p>	<b>(2)</b>

Question Number	Acceptable Answer	Additional Guidance	Mark
9(e)(ii)	<p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>ethanol forms hydrogen bonds (with water) (1)</li> <li>chloroethane forms (permanent) dipole-dipole attractions <b>and</b> London forces (with water) (1)</li> </ul>	<p>Allow "London forces and dipole forces"</p> <p>Ignore 'chloroethane does not form hydrogen bonds with water</p>	(2)

Question Number	Acceptable Answer	Additional Guidance	Mark
9(f)	<ul style="list-style-type: none"> <li>calculation of mol of <math>\text{ClO}^-</math> in 1 000 <math>\text{dm}^3</math> of water (1)</li> <li>calculation of mol of <math>\text{Ca}(\text{ClO})_2</math> (1)</li> <li>calculation of mass of <math>\text{Ca}(\text{ClO})_2</math> (1)</li> </ul>	<p><u>Example of calculation:</u></p> <p><math>5.6 \times 10^{-6} \times 10^3 = 5.6 \times 10^{-3}</math> (mol)</p> <p><math>\frac{5.6 \times 10^{-3}}{2}</math> (mol) = <math>2.8 \times 10^{-3}</math> (mol)</p> <p><math>2.8 \times 10^{-3} \times 143.1 = 0.40068</math> (g) = 0.4 (g)</p> <p>Allow use of 143 instead of 143.1 in M3</p> <p>Ignore SF</p> <p>Answer of 0.8(01) (g) scores (2) marks</p>	(3)

(Total for Question 9 = 23 marks)  
Total for Question Paper = 80 marks

