

Surname	Centre Number	Candidate Number
Other Names		2



GCE AS – NEW AS

B400U10-1



BIOLOGY – Component 1
Basic Biochemistry and Cell Organisation

P.M. THURSDAY, 26 May 2016

1 hour 30 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	9	
2.	6	
3.	7	
4.	16	
5.	13	
6.	15	
7.	9	
Total	75	

ADDITIONAL MATERIALS

In addition to this examination paper, you will need a calculator and a ruler.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation pages at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The assessment of the quality of extended response (QER) will take place in question 7.

The quality of written communication will affect the awarding of marks.




MAY16B400U10101

Answer all questions.

1. Nucleotides are the building blocks of nucleic acids. They have an important role in many biochemical reactions in cell metabolism and regulation, including transport across cell membranes.

The diagram below shows a nucleotide.



- (a) (i) Identify components **A**, **B** and **C**. [1]

A

B

C

- (ii) With reference to components **A** and **B** shown above, describe the composition of the nucleotides of DNA, RNA and ATP. [3]

DNA

.....

RNA

.....

ATP

.....



- (b) An experiment was carried out to investigate the rates of uptake of different sugars by the small intestine. One experiment used normal intestine, the second used a piece of intestine treated with cyanide. The results are shown in the table below.

Sugar	Relative rates of absorption/a.u.	
	Normal intestine	Intestine treated with cyanide
glucose	1.00	0.33
galactose	1.10	0.53
xylose	0.30	0.31
arabinose	0.29	0.29

- (i) Name **two** sugars from the table which can be absorbed by active transport. [1]

.....

- (ii) Using evidence from the table, explain why you chose these sugars. [2]


.....

- (c) It was concluded that all of the sugars named in the table can be absorbed by diffusion. Under what conditions could this take place and how does evidence from the table support this conclusion? [2]


.....



2. Ebola virus disease is a serious illness of humans that originated in Africa, where there was an outbreak which started in 2014. The photographs and diagrams below show the Ebola virus and an enlarged section of the virus in detail.



Ebola virus



Enlarged section of Ebola virus

In addition to the usual viral structure, Ebola is surrounded by a lipid-bilayer, which is derived from infected cell membranes as the virus buds from the cell.

A viral transmembrane glycoprotein, is incorporated into this membrane and allows the virus to bind to blood vessel cells.

- (a) What are the **two** major biochemical components present in **all** viruses? [1]

.....




(b) Describe the biochemical structure of a glycoprotein. [2]

.....

.....

.....

The initial stage of infection involves the following process.



(c) Using information from the diagrams, explain how Ebola virus particles enter a host cell. [3]

.....

.....

.....

.....

.....

.....


.....

.....

B400U101
05



3. (a) Glucose concentration in body fluids can be detected by using a number of tests. In one technique, known concentrations of glucose and reagents are prepared leading to a colour change. The absorbance is then measured using a colorimeter. The results are used to plot a standard (calibration) curve as shown below. Unknown solutions can then be compared with the standard curve to determine their concentration.



Glucose does not appear in urine until blood glucose concentrations of 0.18 mg cm^{-3} and above are reached. This occurs in untreated diabetics.

- (i) When tested, a urine sample gave an absorbance of 0.17 a.u.. State and explain what diagnosis you could reach based on this reading. [2]

.....

.....

.....

.....

- (ii) Why is it not possible to use the standard curve for glucose determination in a **whole blood** sample? [1]

.....

.....



- (b) Modern medicine uses immobilised enzymes, in devices called biosensors, to detect blood glucose levels. This involves the use of glucose oxidase as the enzyme and glucose as the substrate. Give **two** advantages of using immobilised enzymes for blood glucose monitoring. [2]


.....

.....

.....

.....

- (c) The following structures show two monosaccharides.



- (i) Describe the difference between these two molecules. [1]

.....

.....

.....

- (ii) What would be the products of a condensation reaction between these two molecules? [1]

.....



4. An experiment was carried out to determine the change in mass of potato tissue immersed in different sucrose solutions. Skinless potato cylinders were prepared in order to carry out this experiment.

- (a) Calculate the total surface area, to one decimal place, of one of the cylinders which had a length of 45 mm and a diameter of 8 mm. [3]

Surface area of a cylinder is

$$2 \pi r^2 + 2 \pi r h$$

r = radius

h = length

$\pi = 3.14$

Surface area = mm²

Sucrose solutions were prepared at different concentrations. The skinless potato cylinders of equal length from the same potato were weighed and each immersed in one of the solutions. After two hours they were blotted dry and reweighed. The percentage change in mass was calculated.

The results are shown below.


Sucrose Concentration /M	Initial Mass /g	Final Mass /g	Change in Mass /g	Percentage Change in Mass/%
0.0	3.4	4.1	0.7	20.6
0.2	3.4	3.5	0.1	2.9
0.4	6.2	5.3	-0.9	-14.5
0.6	6.3	4.8	-1.5	-23.8
0.8	6.2	4.4	-1.8	-29.0
1.0	6.0	4.3	-1.7	-28.3

- (b) Use the data above to **draw a graph** on the page opposite showing how percentage change in mass of potato is affected by the change in sucrose concentration. [3]



Examiner
only

B400U101
09



- (c) Using the conversion table below, determine the water potential of the potato tissue in this experiment. Explain how you reached this conclusion. [3]

Sucrose concentration /M	Water potential /kPa
0.05	-130
0.10	-260
0.15	-410
0.20	-540
0.25	-680
0.30	-860
0.35	-970
0.40	-1120
0.45	-1280
0.50	-1450
0.55	-1620
0.60	-1800
0.65	-1980
0.70	-2180

.....

.....

.....

.....

.....



(d) Explain what has caused the percentage difference in the mass of the tissue in the following concentrations of sucrose solution in terms of water potential. [4]

0.0 M

.....

.....

0.8 M

.....

.....

(e) The experiment was repeated using the same range of sucrose concentrations and sweet potato but there was no decrease in mass at any sucrose concentration.

(i) Explain why there was no decrease in mass at any of the sucrose concentrations. [1]

.....

.....

(ii) Describe how you would modify the method to determine the ψ_{tissue} of sweet potato. [2]

.....

.....

.....

B400U101
11

16



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**




5. An experiment was set up as shown below with the addition of alkaline sodium carbonate causing the milk to turn pink. Phenolphthalein is a pH indicator which is pink in alkaline conditions and colourless when in acid conditions. The experiment was timed to see how long it took for the indicator to turn from pink to colourless.

stirring rod



add in turn:
5 cm³ milk
7 cm³ sodium carbonate
solution
5 drops of phenolphthalein

stirring rod



1 cm³ lipase solution

add in turn:
5 cm³ milk
7 cm³ sodium carbonate
solution
5 drops of phenolphthalein

stir and start timing when
you add the lipase

- (a) Explain what caused the indicator to lose the pink colour and become colourless. [3]

.....

.....

.....

.....

.....

.....

- (b) Why would this experiment not be suitable for assessing the effect of different pH values on lipase? [1]

.....

.....

.....



Weight loss in obese patients can be achieved by a reduction in their intake of lipids or by reducing their ability to digest and / or absorb lipids.

One drug used to treat obesity acts as a competitive inhibitor of lipase. It prevents approximately 30% of the lipid being absorbed, thereby reducing energy (calorie) intake. The extra lipid in the large intestine can lead to unpleasant side effects such as diarrhoea and deficiency diseases. Treatment of patients requires them to take vitamin tablets since absorption of fat-soluble vitamins and other fat-soluble nutrients is inhibited by the use of the drug. After drug treatment was stopped, a significant number of subjects regained up to 35% of the weight they had lost.

(c) Describe and explain the action of this drug on pancreatic lipase. [3]

.....

.....

.....

.....

.....

.....

(d) (i) Describe how this drug would lead to weight loss. [2]

.....

.....

.....

.....

(ii) With reference to the action of this drug, explain why people tended to regain the weight when they stopped taking it. [1]


.....

.....

.....



During clinical trials of this drug the following results were obtained. Both groups were on weight loss diets but only one group was treated with the drug.



(iii) Using all the information given in (c) and (d), state the advantages and disadvantages of using this drug as part of a weight loss programme. [3]

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



6. In most eukaryotic genes, coding regions (exons) are interrupted by non-coding regions (introns). During transcription, the entire gene is copied into a pre-mRNA molecule, which includes exons and introns. During the process of RNA splicing, introns are removed and exons joined to form a continuous coding sequence. This “mature” mRNA is ready for translation.



(a) Explain the process of transcription including the roles of the enzymes involved. [3]

.....

.....

.....

.....

.....

.....

.....

(b) (i) Explain why it is important that the RNA which is finally translated, consists only of exons. [2]

.....

.....

.....

.....

.....

(ii) Explain how a change in the sequence of bases in the DNA of the gene (a mutation) would affect the protein produced if it was in

I. an intron; [1]

.....

.....

.....

II. an exon. [1]

.....

.....

.....



- (c) (i) The following figures show the mRNA transcribed from two short lengths of DNA. Using the information following, **complete the correct sequence of amino acids** for each mRNA molecule. [1]

Normal

	T	A	C	A	A	A	G	T	C	A	C	C	A	C	T	template strand
	A	U	G	U	U	U	C	A	G	U	G	G	U	G	A	mRNA

Amino acid sequence

Mutation

	T	A	C	A	A	G	G	T	C	A	C	C	A	C	T	template strand
	A	U	G	U	U	C	C	A	G	U	G	G	U	G	A	mRNA

Amino acid sequence

Codons Found in Messenger RNA

Second Base

		<i>Second Base</i>				
		U	C	A	G	
<i>First Base</i>	U	Phe	Ser	Tyr	Cys	U
		Phe	Ser	Tyr	Cys	C
		Leu	Ser	Stop	Stop	A
		Leu	Ser	Stop	Trp	G
	C	Leu	Pro	Hls	Arg	U
		Leu	Pro	Hls	Arg	C
		Leu	Pro	Gln	Arg	A
		Leu	Pro	Gln	Arg	G
	A	Ile	Thr	Asn	Ser	U
		Ile	Thr	Asn	Ser	C
		Ile	Thr	Lys	Arg	A
		Met	Thr	Lys	Arg	G
	G	Val	Ala	Asp	Gly	U
		Val	Ala	Asp	Gly	C
		Val	Ala	Glu	Gly	A
		Val	Ala	Glu	Gly	G



- (ii) Explain why the change of bases shown is described as a silent mutation. [1]

.....

.....

.....

Chargaff carried out work examining the ratios of different bases in samples of DNA from different sources. A sample of his results is shown below.

Percentage composition

Source of DNA	Adenine	Thymine	Guanine	Cytosine
Yeast	31.3	32.9	18.7	17.1
Herring sperm	27.8	27.5	22.1	22.6
Human sperm	30.7	31.2	19.3	18.8


- (d) (i) Calculate the exact ratio to two decimal places of adenine to thymine and guanine to cytosine for human sperm. [2]

adenine to thymine ratio =

guanine to cytosine ratio =



In 1952 Linus Pauling and his co-worker were researching DNA structure and suggested the following structure.




Pauling stated that:

"it is unlikely that the sugar groups constitute the core of the molecule as no satisfactory way of packing them has been found. We conclude that the core of the molecule is probably formed of the phosphate groups arranged in a three chained helix. The purine and pyrimidine groups are on the periphery of the molecule."



At this time, Rosalind Franklin was carrying out X-Ray crystallography on pure DNA samples.



X-ray diffraction image of the double helix structure of the DNA molecule, taken in 1952 by Raymond Gosling, commonly referred to as "Photo 51", during work by Rosalind Franklin on the structure of DNA.

In 1952 **Franklin** concluded that:

"The results suggest a helical structure of DNA, containing probably 2, 3, or 4 nucleic acid chains and having the phosphate groups near the outside."

- (ii) Evaluate how closely the suggestions by **Pauling** and **Franklin**, shown in the boxes above and opposite, **match the current understanding** of the arrangement of nucleotides in DNA. [4]

Pauling

.....

.....

.....

Franklin

.....



.....

.....



7. In multicellular animals cell division is essential for the survival of an individual and its species.

The diagrams below show two types of cell division in the fruit fly *Drosophila*. Discuss the significance of each type of cell division and explain why the cells produced by the process shown in diagram **B** could lead to the formation of a tumour whilst those produced by diagram **A** do not. [9]



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Examiner
only

A large rectangular area with a solid top and bottom border and a dotted line for a left margin, providing space for writing.



A large rectangular area with a solid top and bottom border and a solid left and right border. Inside this area, there are 24 horizontal dotted lines spaced evenly down the page, providing a writing area for the student.



Area with horizontal dotted lines for writing.

END OF PAPER

9



BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**



