



GCE AS MARKING SCHEME

SUMMER 2019

**AS
CHEMISTRY - COMPONENT 1
B410U10-1**

INTRODUCTION

This marking scheme was used by WJEC for the 2019 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

COMPONENT 1: THE LANGUAGE OF CHEMISTRY, STRUCTURE OF MATTER AND SIMPLE REACTIONS

SUMMER 2019 MARK SCHEME

GENERAL INSTRUCTIONS

Recording of marks

Examiners must mark in red ink.

One tick must equate to one mark, apart from extended response questions where a level of response mark scheme is applied.

Question totals should be written in the box at the end of the question.

Question totals should be entered onto the grid on the front cover and these should be added to give the script total for each candidate.

Extended response questions

A level of response mark scheme is applied. The complete response should be read in order to establish the most appropriate band. Award the higher mark if there is a good match with content and communication criteria. Award the lower mark if either content or communication barely meets the criteria.

Marking rules

All work should be seen to have been marked.

Marking schemes will indicate when explicit working is deemed to be a necessary part of a correct answer.

Crossed out responses not replaced should be marked.

Marking abbreviations


The following may be used in marking schemes or in the marking of scripts to indicate reasons for the marks awarded.

cao = correct answer only
ecf = error carried forward
bod = benefit of doubt

Credit should be awarded for correct and relevant alternative responses which are not recorded in the mark scheme.

Section A

Question				Marking details	Marks available					
					AO1	AO2	AO3	Total	Maths	Prac
1				$3p^64s^23d^{10}4p^6$		1		1		
2				$^{28}_{14}\text{Si}$		1		1		
3	(a)			(molecules are) polar if their atoms have different electronegativities / different tendencies to attract electrons in a bonded pair	1			1		
	(b)			$\begin{array}{cc} & \delta- \quad \delta+ \\ \text{F}-\text{F} & \text{F}-\text{Cl} \end{array}$	1			1		
4				Halogen chlorine Use (water) sterilisation (1) both needed Halide fluoride Use in toothpaste / water supply to prevent tooth decay (1) both needed				2		
5				any value > 500 but < 1000			1	1		


Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
6			diagram to show face centred cubic structure of Na ⁺ and Cl ⁻ (ions on minimum of 2 faces) e.g. <div style="text-align: center;">  </div>	1			1		
7			$K_c = \frac{[AB_2]}{[A][B]^2} \quad (1)$ $1.47 = \frac{0.4}{0.2[B]^2}$ $[B] = 1.17 \text{ (mol dm}^{-3}\text{)} \quad (1)$ no ECF from incorrect K_c expression		2		2	2	
Section A total				5	4	1	10	2	0

Section B

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
8	(a)		<p>the Periodic Table has the elements arranged by increasing number of protons (1)</p> <p>the (atomic) masses are due to protons and neutrons (1)</p> <p>in argon there are more atoms with greater number of neutrons (1)</p>			3	3		


Question		Marking details	Marks available					
			AO1	AO2	AO3	Total	Maths	Prac
(b)	(i)	<p>Indicative content</p> <ul style="list-style-type: none"> • element is vaporised • ionised by having electron knocked off / electron gun • forms positive ions • accelerated by charged plates • made into stream by slits • deflected by electromagnet • degree of deflection according to m/z • detected • range of magnetic fields • whole system under vacuum <p>5-6 marks All stages mentioned with some explanation of what happens at each stage and where each stage occurs <i>The candidate constructs a relevant, coherent and logically structured method including all key elements of the indicative content. A sustained and substantiated line of reasoning is evident and scientific conventions and vocabulary is used accurately throughout.</i></p> <p>3-4 marks Most stages mentioned with some attempt at explaining what happens at each stage or where each stage occurs <i>The candidate constructs a coherent account including most of the key elements of the indicative content. Some reasoning is evident in the linking of key points and use of scientific conventions and vocabulary are generally sound.</i></p> <p>1-2 marks Some stages mentioned with limited description of what happens where <i>The candidate attempts to link at least two relevant points from the indicative content. Coherence is limited by omission and/or inclusion of irrelevant material. There is some evidence of appropriate use of scientific conventions and vocabulary.</i></p> <p>0 marks <i>The candidate does not make any attempt or give an answer worthy of credit.</i></p>	6			6		

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
		(ii)	$\frac{(15 \times 20) + (5 \times 21)}{20} \quad (1)$ <p>accept any correct abundance ratio</p> $20.25 / 20.3 \quad (1)$		2		2	2	
		(iii)	$^{20}\text{T}^{2+}$			1	1		
			Question 8 total	6	2	4	12	2	0

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
9	(a)		$X(g) \rightarrow X^+(g) + e$ must include state symbols	1			1		
	(b)	(i)	increasing number of protons present (in the nucleus) / greater nuclear charge (1) greater attraction therefore more energy is needed to remove electron (1) ignore references to shielding	2			2		
		(ii)	Group 5 (1) (small) fall to the next element / Group 6 element as a paired electron is being removed (1) do not award explanation mark if incorrect group given		2		2		
		(iii)	Z marked below level of all other points e.g. 			1	1		

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
	(c)	(i)	award (1) for either of following <ul style="list-style-type: none"> the number of particles in 1 mol the number of ^{12}C atoms in 12 g of carbon-12 	1			1		
		(ii)	34.23 g is 0.1 mol (1) this has 6.02×10^{22} particles (1) each $\text{Al}_2(\text{SO}_4)_3$ contains 12 oxygen atoms therefore number of oxygen atoms is 7.22×10^{23} (1)		3		3	3	
			Question 9 total	4	5	1	10	3	0

Question		Marking details		Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
10	(a)		<p>precipitate of barium sulfate forms with barium chloride (1)</p> <p>none with magnesium chloride as magnesium sulfate is appreciably soluble (1)</p> <p>award (1) for either of following equations</p> $\text{Ba}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s})$ $\text{BaCl}_2(\text{aq}) + \text{H}_2\text{SO}_4(\text{aq}) \rightarrow \text{BaSO}_4(\text{s}) + 2\text{HCl}(\text{aq})$		3		3		3
	(b)		<p>there are 6 bond pairs (1)</p> <p>these repel to maximum separation / minimum repulsion (1)</p> <p>award (1) for 3D diagram to show octahedral structure e.g.</p> <div style="text-align: center;"> </div>	3			3		
	(c)		<p>electrons are excited by electricity (1)</p> <p>promoted to a higher energy level (1)</p> <p>then fall back (to lower level) and give out energy (1)</p> <p>the energy emitted is in the yellow part of the visible spectrum (1)</p>	4			4		
		Question 10 total		7	3	0	10	0	3

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
11	(a)	(i)	award (1) for four bonding pairs award (1) for octet around all chlorine atoms e.g. 		2		2		
		(ii)	109° / 109.5°	1			1		
		(iii)	$\text{SiCl}_4(\text{l}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{SiO}_2(\text{s}) + 4\text{HCl}(\text{aq})$ award (1) for all formulae correct award (1) for correct balancing and state symbols			2	2		2
		(iv)	SiCl ₄ is a simple (covalent) molecule (1) SiO ₂ is a giant (covalent) molecule (1) SiCl ₄ has weak intermolecular forces (1) strong covalent bonds need to be broken to melt SiO ₂ (1)	4			4		

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
(b)	(i)		pH = $-\log [\text{H}^+]$ (1) 0.22 (1)	1			2	2	
	(ii)		1.26×10^{-7}		1		1	1	
	(iii)		lower pH means that $[\text{H}^+]$ has gone up / is higher than expected (1) equilibrium has moved to RHS (1) forward reaction must be endothermic (1)			3	3		
(c)			theoretical yield = $\frac{5}{24.3} \times 22.4 = 4.61$ (1) percentage purity = $\frac{4.31}{4.61} \times 100 = 93.5\%$ (1) accept alternative method award (1) for actual or theoretical number of moles of hydrogen <ul style="list-style-type: none"> actual $n(\text{H}_2) = \frac{4.31}{22.4} = 0.192$ mol theoretical $n(\text{H}_2) = \frac{5}{24.3} = 0.206$ mol percentage purity = $\frac{0.192}{0.206} \times 100 = 93.2\%$ (1)		2		2	2	2
Question 11 total				6	6	5	17	5	4

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
12	(a)	(i)	(+)5		1		1		
		(ii)	$n(\text{NaClO}_3) = 0.826$ (1) $n(\text{O}_2) = 0.826 \times 1.50 = 1.239$ (1) $V = \frac{nRT}{p} = \frac{1.239 \times 8.31 \times 600}{1.01 \times 10^5} = 61.2 \text{ dm}^3$ (1) must be to 3 sig figs accept molar volume method $V = \frac{1.239 \times 22.4 \times 600}{273} = 61.0 \text{ dm}^3$ (1) must be to 3 sig figs		2	1	3	3	
	(b)		$M_r(\text{NaClO}) = 74.5$ total M_r of reactants = 151.02 (1) both needed $\text{atom economy} = \frac{74.5}{151.02} \times 100 = 49.3\%$ (1)		2		2	1	
	(c)		$\frac{18.8}{23} : \frac{29.0}{35.5} : \frac{52.2}{16}$ $0.817 : 0.817 : 3.27$ (1) $1 : 1 : 4$ therefore empirical formula is NaClO_4 (1)		2		2	1	
Question 12 total				0	7	1	8	5	0

Question			Marking details	Marks available						
				AO1	AO2	AO3	Total	Maths	Prac	
13	(a)		to remove insoluble / solid impurities	1			1		1	
	(b)		(250cm ³) volumetric flask	1			1		1	
	(c)		to show the end-point	1			1		1	
	(d)		as many as needed until two/three concordant titres are obtained (to ensure correct answer)	1			1		1	
	(e)		BaCO ₃ (s) + 2HCl(aq) → BaCl ₂ (aq) + CO ₂ (g) + H ₂ O(l)		1		1			
	(f)		$n(\text{HCl}) = \frac{250}{1000} \times 0.5 = 0.125 \text{ mol}$		1		1			
	(g)		number of moles HCl neutralised in each titration $= 27.8 \times \frac{0.1}{1000} = 2.78 \times 10^{-3} \text{ mol} \quad (1)$ number of moles HCl left after reacting with ore $= 2.78 \times 10^{-3} \times 10 = 2.78 \times 10^{-2} \text{ mol} \quad (1)$ therefore number of moles neutralised by ore $= 0.125 - 2.78 \times 10^{-2} = 0.0972 \text{ mol} \quad (1)$				3	3	3	3
	(h)		number of moles BaCO ₃ = $\frac{0.0972}{2} = 0.0486 \text{ mol} \quad (1)$ mass barium in ore = $0.0486 \times 137 = 6.66 \text{ g} \quad (1)$ percentage barium = $\frac{6.66}{19.15} \times 100 = 34.8 \% \quad (1)$		3		3	3	3	

Question				Marking details	Marks available					
					AO1	AO2	AO3	Total	Maths	Prac
	(i)			insoluble barium compound (e.g. barium sulfate) could have been present			1	1		1
				Question 13 total	4	5	4	13	6	11

COMPONENT 1: THE LANGUAGE OF CHEMISTRY, STRUCTURE OF MATTER AND SIMPLE REACTIONS

SUMMARY OF MARKS ALLOCATED TO ASSESSMENT OBJECTIVES

Question	AO1	AO2	AO3	Total	Maths	Prac
Section A	5	4	1	10	2	0
8	6	2	4	12	2	0
9	4	5	1	10	3	0
10	7	3	0	10	0	3
11	6	6	5	17	5	4
12	0	7	1	8	5	0
13	4	5	4	13	6	11
Totals	32	32	16	80	23	18