

# A Level Chemistry B (Salters) H433/01 Fundamentals of chemistry

Sample Question Paper

# Date - Morning/Afternoon

Time allowed: 2 hours 15 minutes



• the Data Sheet for Chemistry B (Salters)

#### You may use:

· a scientific calculator



First name	
Last name	
Centre number	Candidate number

#### **INSTRUCTIONS**

- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer all the questions.
- · Write your answer to each question in the space provided.
- Additional paper may be used if required but you must clearly show your candidate number, centre number and question number(s).
- Do not write in the barcodes.

#### **INFORMATION**

- The total mark for this paper is **110**.
- The marks for each question are shown in brackets [ ].
- Quality of extended responses will be assessed in questions marked with an asterisk (\*).
- This document consists of 36 pages.



# **SECTION A**


# You should spend a maximum of 40 minutes on this section.

Answer **all** the questions.

1	Whic	h two compounds can react together to form a condensation polymer?	
	A	$C_2H_4$ and $C_3H_6$	
	В	C <sub>2</sub> H <sub>5</sub> OH and CH <sub>3</sub> COOH	
	C	H <sub>2</sub> N(CH <sub>2</sub> ) <sub>6</sub> NH <sub>2</sub> and CH <sub>3</sub> COOH	
	D	HOOC(CH <sub>2</sub> ) <sub>4</sub> COOH and HOCH <sub>2</sub> CH <sub>2</sub> OH	
	Your	answer	[1]
2	Whic	h of the following is a correct statement about the test for Fe <sup>2+</sup> ions in solution?	
	A	add NaOH(aq); brown precipitate formed, soluble in NH <sub>3</sub> (aq)	
	В	add NaOH(aq); brown precipitate formed, insoluble in NH <sub>3</sub> (aq)	
	C	add NaOH(aq); green precipitate formed, soluble in NH <sub>3</sub> (aq)	
	D	add NaOH(aq); green precipitate formed, insoluble in NH <sub>3</sub> (aq)	
	Your	answer	[1]

3 Instrumental techniques are crucially important in the analysis of many mixtures. These may involve wide ranging areas such as in forensic work or the analysis of a potential forged painting.

The figure below shows a graphical analysis of a mixture using a particular technique.



Identify the technique used to produce this graph.

- **A** mass spectrometry
- **B** infrared spectroscopy
- C proton NMR spectroscopy
- **D** gas-liquid chromatography

Your answer	
Your answer	

[1]

- **4** Which of the following factors is most important in determining the relative reactivity of the haloalkanes?
  - **A** bond polarity
  - **B** bond enthalpy
  - **C** intermolecular bonding
  - **D** molar mass

Your answer	

[1]

5	Choo	se the statement that is <b>not</b> a principle of 'green chemistry'.	
	A	Cleaning up waste after it is formed.	
	В	Keeping the atom economy as high as possible.	
	C	Use of selective catalysts.	
	D	Choosing reactants to avoid explosions.	
	Your	answer	[1]
6		emplete combustion, a sample of a hydrocarbon produces 33 g of carbon dioxide and of water.	
	What	is the empirical formula of this hydrocarbon?	
	A	$C_3H_8$	
	В	$CH_2$	
	C	$C_2H_4$	
	D	$C_3H_4$	
	Your	answer	[1]
7	38.49	g of MgSO <sub>4</sub> •xH <sub>2</sub> O crystals were heated to drive off the water. 18.79 g of solid remained.	
	What	is the value of $x$ ?	
	A	1	
	В	2	
	C	6	
	D	7	
	Your	answer	[1]

8	Stront	tium oxide reacts with water to form an alkaline solution.	
	What	mass of water would react with 10 g strontium oxide?	
	A	0.94 g	
	В	1.7 g	
	C	1.9 g	
	D	3.5 g	
	Your	answer	[1]
9		mass of sodium hydroxide should be dissolved in 250 cm <sup>3</sup> of water to make a solution that concentration of 0.05 mol dm <sup>-3</sup> ?	
	A	0.0125 g	
	В	0.5 g	
	C	2 g	
	D	8 g	
	Your	answer	[1]
10	Which	n of the following is a correct statement about the bonding in a CO molecule?	
	A	Carbon provides an electron pair in a dative covalent bond.	
	В	Oxygen has two lone pairs of electrons.	
	C	There are three covalent bonds between the atoms.	
	D	The molecule has no dipole.	
	Your	answer	[1]

		O	
Which molecule has	the smallest bo	ond angle?	
A BCl <sub>3</sub>			
$\mathbf{B}$ BeCl <sub>2</sub>			
C NCl <sub>3</sub>			
<b>D</b> SiCl <sub>4</sub>			
Your answer			[1]
		<b>Z</b> . Each contains one t is not known which	e of potassium chloride, potassium solution is which.
-		•	ach of <b>X</b> , <b>Y</b> and <b>Z</b> . In each case the student cyclohexane layer are shown below.
Γ	Solution X	red/brown colour	
	Solution Y	colourless	
-	Solution Z	violet colour	
<ul> <li>A X = potassium</li> <li>B X = potassium</li> <li>C X = potassium</li> </ul>	m chloride, <b>Y</b> = m bromide, <b>Y</b> = m chloride, <b>Y</b> =	ties of solutions <b>X</b> , <b>Y</b> potassium bromide,  potassium chloride,  potassium iodide, <b>Z</b> potassium chloride, <b>Z</b>	<ul><li>Z = potassium iodide</li><li>Z = potassium iodide</li><li>= potassium bromide</li></ul>
Your answer			[1]

A teacher tests a solution that has been used by students in some experiments with potassium halides to see if it can be used with a different class.

The teacher adds silver nitrate solution to the solution that the students were using. A pale yellow precipitate is formed. The teacher adds a few drops of dilute ammonia solution to the precipitate. Some of the precipitate appears to dissolve and the precipitate remaining has a stronger yellow tinge. The teacher adds excess concentrated ammonia solution to the remaining precipitate. The precipitate does not dissolve.

What is the most likely conclusion that the teacher will reach about the solution that the students were using?

		6	
	A	The solution contains chloride ions.	
	В	The solution contains bromide ions.	
	C	The solution contains chloride and bromide ions.	
	D	The solution contains chloride and iodide ions.	
	Your	answer	[1]
14		gases, $W$ , $X$ , $Y$ and $Z$ , are known to be $N_2$ , $NO$ , $NH_3$ and $O_2$ .	
	It is 1	not known which gas is which.	
		n gases <b>X</b> and <b>Y</b> are mixed a brown gas is formed.	
		Y relights a glowing splint.	
		s W, X and Z extinguish a burning splint. Z turns red litmus paper blue.	
	Gas A	z turns red nunus paper blue.	
	Whic	ch gas is N <sub>2</sub> ?	
	A	Gas W	
	В	Gas <b>X</b>	
	C	Gas Y	
	D	Gas <b>Z</b>	
	Your	answer	[1]

15	A wh	ite solid has the following properties:	
	It giv	melts at the same temperature. es a purple colour with neutral FeCl <sub>3</sub> (aq). e is fizzing when Na <sub>2</sub> CO <sub>3</sub> (aq) is added.	
	The v	white solid could be	
	A	A mixture of a phenol and a carboxylic acid.	
	В	A compound with phenol and carboxylic acid functional groups.	
	C	A phenol.	
	D	A carboxylic acid.	
	Your	answer	[1]
16	RNA and b	is a polymer made of nucleotides. Nucleotides are monomers made of phosphates, sugars, ases.	1
	Whic	h statement is correct about RNA?	
	A	Phosphate and deoxyribose units join together by a condensation reaction to form the phosphate–sugar backbone in RNA.	
	В	In RNA, adenine forms a base pair with thymine; cytosine forms a base pair with guanin	e.
	C	Bases join to the phosphate–sugar backbone by a condensation reaction.	
	D	Base pairs form by a condensation reaction between the bases.	
	Your	answer	[1]

17	The process	of protein	synthesis	involves	DNA	and RNA	in the	following	steps.
----	-------------	------------	-----------	----------	-----	---------	--------	-----------	--------

- 1. An anti-codon on tRNA binds to a triplet code on mRNA.
- 2. A ribosome moves along a strand of mRNA.
- 3. DNA transcription occurs to make a strand of mRNA.
- **4.** Amino acids carried by tRNA are assembled into a protein chain.

What is the correct order of these steps in protein synthesis?

- **A** 3-2-1-4
- **B** 3-1-2-4
- $\mathbf{C}$  2 3 4 1
- **D** 1-3-4-2

Your answer	

[1]

18 What template DNA sequence will produce the polypeptide GluValLeu?


You may need to refer to your Data Sheet.

- A CUGGUCGAA
- **B** GTTCAGGAC
- C GUUCAGGAC
- **D** CAGGACTTC

Your	answer	

[1]

19 The diagram below shows the structure of a penicillin. The properties of the penicillin can be modified by changing the 'R' group side chain.




What name is given to the part of the drug that is medicinally active?

- A modifier
- **B** pharmacophore
- C receptor
- **D** chromophore

Your answer

[1]

**20** Which one of the following graphs correctly represents the shape of a rate versus substrate concentration plot for an enzyme-catalysed reaction?



21 Aspartame is a sweetener which can be used instead of sugar. It is a methyl ester of a dipeptide formed from a reaction between the amino acids aspartic acid and phenylalanine, the structures of which are shown below.

$$\begin{array}{c} \text{NH}_2 \\ \text{OH} \\ \text{OH} \end{array}$$

Aspartic acid

Phenylalanine

Alkaline hydrolysis of aspartame produces salts of the above amino acids.

Which pair of compounds is produced?

**A**


$$Na^{+}O^{-} = 0$$

$$ONa^{+} = 0$$

© OCR 2014 H433/01

Your answer

22 Below is an enthalpy-level diagram for the solution of lithium bromide.



Which enthalpy change is represented by  $\Delta H_2$ ?

- $\mathbf{A}$   $\Delta_{\text{sol}}H(\text{Li}^+) + \Delta_{\text{sol}}H(\text{Br}^-)$
- $\mathbf{B}$   $\Delta_{\text{hyd}}H(\text{Li}^+) + \Delta_{\text{hyd}}H(\text{Br}^-)$
- $\mathbf{C}$   $\Delta_{\mathrm{LE}}H(\mathrm{LiBr})$
- $\mathbf{D}$   $\Delta_{\text{sol}}H(\text{LiBr})$

Your answer

[1]

23 Four changes W, X, Y and Z have the following signs for their  $\Delta H$  and  $\Delta_{sys}S$  values.

	W	X	Y	Z
$\Delta H$	+	+	_	_
$\Delta_{ m sys}S$	+	1	+	_

Which of the following is a correct statement?

- A Change W never occurs.
- **B** Change **X** could be the reaction of a solid carbonate with an acid.
- C Change Y could be a precipitation reaction.
- **D** Change **Z** could be the condensation of a gas to a liquid.

Your answer		[1]

- 24 A pure sample of fat was known to be a triester of propane-1,2,3-triol with a fatty acid.
  - 0.15 mol of the fat required  $10.8 \text{ dm}^3$  of hydrogen at room temperature and pressure for complete hydrogenation.

Identify which fatty acid formed the triester.

- **A** Palmitic acid, C<sub>15</sub>H<sub>31</sub>COOH
- **B** Linoleic acid, C<sub>17</sub>H<sub>31</sub>COOH
- C Oleic acid, C<sub>17</sub>H<sub>33</sub>COOH
- **D** Stearic acid, C<sub>17</sub>H<sub>35</sub>COOH

Your answer			

[1]

25 Chemists use knowledge of bonding between molecules and ions to explain why different substances are soluble in water.

Which of the following statements is correct?

- **A** pentane is insoluble in water because pentane molecules are non-polar and **do not** form hydrogen bonds with water molecules
- **B** propanone is soluble in water even though water molecules **do not** form hydrogen bonds with propanone molecules
- C sodium chloride is soluble in water because water molecules are polar and form hydrogen bonds with Na<sup>+</sup> ions and Cl<sup>-</sup> ions
- **D** carbon dioxide is insoluble in water because it is non-polar and **does not** form hydrogen bonds with water

Your answer	

[1]

26 The decomposition of calcium carbonate is a reaction that takes place in cement manufacture.

$$CaCO_3 \rightarrow CaO + CO_2$$

$$\Delta_{\rm f}H = +178 \text{ kJ mol}^{-1}$$

	CaCO <sub>3</sub>	CaO	$CO_2$
$S / J \text{ mol}^{-1} K^{-1}$	93	40	214

What is  $\Delta_{tot}S$  for this reaction at 298 K?

**A** 
$$-436 \text{ J mol}^{-1} \text{ K}^{-1}$$

**B** 
$$+160 \text{ J mol}^{-1} \text{ K}^{-1}$$

$$\mathbf{C}$$
 +161 J mol<sup>-1</sup> K<sup>-1</sup>

$$\mathbf{D}$$
 +758 J mol<sup>-1</sup> K<sup>-1</sup>

Your answer	

[1]

A few drops of universal indicator solution are added to a solution of sodium chloride. Two

Statement 2:	e solution turns red e solution turns colourless	
	e solution turns colourless	
Statement 3: The statement of the statem		
	e solution turns green	
<ul><li>A 1, 2 and 3</li><li>B Only 1 and 2</li></ul>		
C Only 2 and 3		
<b>D</b> Only 1		

28 Some possible ligands are shown below.

Which of the following could be bidentate ligands?

- 1: NH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>
- **2:**  $(COO^{-})_{2}$
- $C_2H_5OH$
- **A** 1, 2 and 3

**27** 

- B Only 1 and 2
- C Only 2 and 3
- **D** Only 1

Your	answer	


[1]

29	Which statement correct?	t(s) about the reaction of aqueous sodium hydroxide with 1-chloropentane is/are	
	<b>Statement 1:</b>	The reaction is a substitution reaction	
	<b>Statement 2:</b>	The reaction occurs at a slower rate than with 1-bromopentane	
	<b>Statement 3:</b>	'Curly arrows' can be used to show the movement of pairs of electrons in the mechanism for this reaction	
	<b>A</b> 1, 2 and 3	3	
	<b>B</b> Only 1 ar	nd 2	
	C Only 2 ar	nd 3	
	<b>D</b> Only 1		
	Your answer		[1]
30		lowing quantities increase from left to right along the series: SrSO <sub>4</sub> ; BaSO <sub>4</sub> ?	
	1:	Size of the metal ions	
	2:	The magnitude of enthalpy change of hydration of the metal ions	
	3:	The magnitude of lattice enthalpy of the compounds	
	<b>A</b> 1, 2 and 3		
	<b>B</b> Only 1 ar	nd 2	
	C Only 2 ar	nd 3	
	<b>D</b> Only 1		
	Your answer		[1]

## **SECTION B**

### Answer **all** the questions.

31 Ceria, CeO<sub>2</sub>, is used in a thermochemical cycle for splitting water. A simplified diagram of the process is shown below.



(a)	(i)	Give the systematic name for ceria.	
			[1]
	(ii)	Give chemical equations for the <b>two</b> reactions involved in the cycle.	
		State symbols are not required.	
		equation 1:	
		equation 2:	[2]
	(iii)	One of the reactions in the cycle is endothermic.	
		What data for each reactant and product is required to decide which reaction is endothermic	?

[1]

- **(b)** Aqueous solutions of  $Ce(SO_4)_2$  and  $KMnO_4$  can be used as oxidising agents in redox titrations. In both cases reactions are carried out in acidic solutions.
  - (i) KMnO<sub>4</sub> should **not** be acidified with hydrochloric acid.

Select and use data from the table below to determine which acid can be used with  $Ce^{4+}$  to oxidise  $Cl^-$ .

Explain your answer.

Half-reaction	$E^{\theta}$ / V
$Ce^{4+} + e^{-} \rightleftharpoons Ce^{3+}$	+1.28 measured in 1 mol dm <sup>-3</sup> HCl(aq)
$Cl_2 + 2e^- \Rightarrow 2Cl^-$	+1.36
$Ce^{4+} + e^{-} \Rightarrow Ce^{3+}$	+1.44 measured in 1 mol dm <sup>-3</sup> H <sub>2</sub> SO <sub>4</sub> (aq)
$MnO_4^- + 8H^+ + 5e^- \Rightarrow Mn^{2+} + 4H_2O$	+1.51


(ii) A student devises an experiment to find out how much cerium is present in an impure sample of ceria.

First the student converts 2.5 g of powdered ceria into  $100 \text{ cm}^3$  of a solution of  $\text{Ce}(\text{SO}_4)_2$  in sulfuric acid. Then 25.0 cm<sup>3</sup> of this  $\text{Ce}(\text{SO}_4)_2$  solution is titrated with a standard solution of arsenic(III) in sulfuric acid. In the reaction  $\text{Ce}^{4+}$  is reduced to  $\text{Ce}^{3+}$ .

 $1.00~\text{dm}^3$  of solution is made up by dissolving 0.250~mol of  $As_2O_3$  in sulfuric acid. The arsenic is present in the solution as  $AsO_3^{3-}$  ions.  $25.0~\text{cm}^3$  samples of the  $Ce(SO_4)_2$  solution required an average titre of  $3.00~\text{cm}^3$  of this  $AsO_3^{3-}$  solution.

The half-equation for the oxidation of AsO<sub>3</sub><sup>3-</sup> ions is given below.

$$AsO_3^{3-} + H_2O \rightarrow AsO_4^{3-} + 2H^+ + 2e^-$$


Calculate the percentage purity of the student's ceria sample.

Show all your working.

Give your answer to an **appropriate** number of significant figures.

(iii) Suggest one improvement to the student's method that will reduce the percentage error in the answer to (ii) without changing the apparatus.

Give a reason for your choice.




(iv)	Calculate the number of oxygen atoms in 2.5 g of <b>pure</b> ceria.
	Show all your working.
	number of atoms = $\dots$ [2]
( <b>v</b> )	Ceria is also used as a heterogeneous oxidation catalyst for self-cleaning ovens. Ceria is incorporated on the surface of the inside panels of an oven.
	Describe how a heterogeneous catalyst works.
	[4]

<b>32</b>	Ethanol is sometimes used as a biofuel to replace petrol in car engines. However it has several
	disadvantages.


(a)	Give two disadvantages of ethanol as a replacement fuel for petrol.			
		• • • • •		
		[2]		
	•••••••••••••••••••••••••••••••	L4J		

**(b)\*** The mass spectrum and <sup>13</sup>C NMR spectrum for compound **A** are given below.

# Mass spectrum (Compound A)



<sup>13</sup>C NMR spectrum (Compound A)



Use the information below and the spectral data given on the previous page to work out the **structural formulae** of compounds **A**, **B**, **C** and **D**.

Compound A reacts when heated with acidified K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> to form compound B.

Compound **A** has the molecular formula  $C_xH_yO$ .

Compound <b>B</b> does <b>not</b> react with Tollens' reagent or NaOH.  Compound <b>A</b> reacts with heated $Al_2O_3$ to form two unsaturated compounds, <b>C</b> and <b>D</b> .
Include evidence to support your choice of structures.
•••••••••••••••••••••••••••••••••••••••
[6]

(c)	The electricity-tolerant bacteria used to convert methanoic acid contain an enzyme used to synthesise compound <b>A</b> . This enzyme is specific to this synthesis.
	Explain why an enzyme is specific for a particular reaction.
	[2]
( <b>d</b> )	A mixture of 4.0 g of ethanol vapour and 25 dm <sup>3</sup> of air is ignited.
	Determine whether the ethanol will be completely burned.
	All volumes are measured at room temperature and pressure.
	Assume air contains 21% oxygen.

## **BLANK PAGE**



33	Coconut oil contains a large amount of compound E. The sodium salt, F, of compound E is used as a
	soap.

#### Compound E

(a)	Circle an ester	group on	the diagram	of the structure	of <b>E</b> above.
-----	-----------------	----------	-------------	------------------	--------------------

[1]

[1]

- **(b)** After **F** is formed from **E**, **F** remains dissolved in the solution. **F** is precipitated out as a solid soap by adding NaCl.
  - (i) Using  $Na^+L^-$  to represent the solid  $\mathbf{F}$ , give the mathematical equation for the solubility product of  $\mathbf{F}$ .

Explain why adding NaCl to a solution of <b>F</b> causes the salt to precipitate out.	
	••••
	 [2]
	Explain why adding NaCl to a solution of <b>F</b> causes the salt to precipitate out.

(iii)	The solubility in water of <b>F</b> is 24.0 g dm <sup><math>-3</math></sup> at 24 °C.
	The molar mass of $\mathbf{F}$ is 222 g mol <sup>-1</sup> .
	Calculate the minimum mass (in g) of solid NaCl that is needed to form a precipitate when added to $500~\text{cm}^3$ of a $1.00\times10^{-2}~\text{mol dm}^{-3}$ of <b>F</b> at 24 °C.
	Assume you can neglect the $[Na^+(aq)]$ from salt ${\bf F}$ in comparison with the $[Na^+(aq)]$ from NaCl.
	Show all your working.
	Give one <b>other</b> assumption you make in your calculation.
	minimum mass of NaCl = g
assu	Imption:
••••	
••••	

[5]

(c) Ester G is a fruit-flavoured additive for some ice cream and chewing gum. Ester G can be hydrolysed to form acid H and methanol.

$$\begin{array}{c|ccccc} CH_2COOCH_3 & & & H^+ & CH_2COOH \\ & & + & 2H_2O & & & \\ CH_2COOCH_3 & & & CH_2COOH \end{array} & + & 2CH_3OH \\ & & & & \\ Ester G & & & Acid H \end{array}$$

A mixture of ester  ${\bf G}$  and water is left to reach equilibrium with a small amount of acid catalyst.

The initial concentrations are:

Ester G: 2.0 mol dm<sup>-3</sup>
 Water: 5.0 mol dm<sup>-3</sup>

The equilibrium concentration of ester G is 1.4 mol dm<sup>-3</sup>.

Calculate the value of  $K_c$  for this equilibrium.

Show all your working.

$$K_{\rm c} = \dots$$
 [4]

(d) Sodium ethanoate may be used in the production of ester **G**. It can also be used in buffer solutions.

Calculate the mass of solid sodium ethanoate (molar mass =  $82 \text{ g mol}^{-1}$ ) that would need to be added to ethanoic acid (concentration  $1.0 \text{ mol dm}^{-3}$ ) in order to make  $250 \text{ cm}^3$  of buffer solution with a pH of 4.80.

 $K_a$  of ethanoic acid is  $1.74 \times 10^{-5}$  mol dm<sup>-3</sup>.

Show all your working.

mass of solid sodium ethanoate = ...... g [4]

34	and	pberry ketone, extracted from various berries, is widely used as a flavouring, a constituent of perfumes as a diet supplement. The cost of natural raspberry ketone is often over 2000 times the cost of berry ketone manufactured synthetically.
	(a)	Suggest a reason why natural raspberry ketone is expensive to produce.
		[1]
	(b)*	The structures of raspberry ketone and cyclohexanol are shown below.
		OH OH
		raspberry ketone cyclohexanol
		The difference between the structures of the carbon ring in raspberry ketone and cyclohexanol has an effect on the chemical properties of the hydroxyl group.
		Describe the difference between the structures of the two carbon rings and how the reactions of the hydroxyl group are different in the two compounds.
		Include examples, and at least two equations.

**[6]** 

<b>(c)</b>	Raspberry ketone reacts with chloromethane in the presence of anhydrous aluminium chloride to
	form a mixture of two structural isomers.

What technique could be used to show that two organic products are present in the mixture?

.....[1]

(d) Cyanide ions will react with raspberry ketone.


Complete the diagram below to show the mechanism for the formation of a cyanhydrin, showing the intermediate and the product.

Include curly arrows and partial charges where appropriate.




[4]

(e) Raspberry ketone can be reduced to the compound shown below.



Explain why this compound has two stereoisomers and describe how they differ.	
•••••••••••••••••••••••••••••••••••	•••••
	•••••
	[2]
	[-]

**(f)** 4-Phenylbutan-2-one is a liquid which boils at 235 °C whereas raspberry ketone is a solid, melting at 83 °C.



4-phenylbutan-2-one

Explain this differe	ence in properties.		
•••••		•••••	 
••••••		••••••	

35	Alu	miniu	m is produced by the electrolysis of molten aluminium oxide, Al <sub>2</sub> O <sub>3</sub> .
	(a)	Den	nonstrate how the formula of aluminium oxide can be deduced using the periodic table.
		••••	
		••••	
		••••	[1
	<b>(b)</b>	(i)	Write the equation for the reaction at the cathode when molten aluminium oxide is electrolysed
			[1
		(ii)	The oxygen produced at the anode reacts with the carbon electrodes.
			Calculate the maximum volume of carbon dioxide (measured at RTP) that would be produced during the electrolysis of $100~\rm kg$ of $Al_2O_3$ .
			Show all your working.
			volume = $dm^3$ [3

(c) (	(i)	ninium is not produced at the cathode when a solution of an aluminium salt is electrolysed.	
		A solution of aluminium sulfate is electrolysed.	
		Suggest equations for:	
		• the reaction at the cathode	
		• the reaction at a carbon anode.	
			[2]
(	(ii)	Give details of a simple test that can be used to test for the sulfate ion in solution.	
			•••
			[2]

(d) Aluminium can be used to clean silver cutlery that has been 'tarnished' by a layer of black silver sulfide. The silver object is placed in a hot solution of sodium carbonate in contact with a piece of aluminium foil. An electrochemical reaction occurs and the tarnished silver becomes shiny again.

	<i>E</i> <sup>⊕</sup> / V
$Al^{3+} + 3e^- \Rightarrow Al$	-1.66
$Ag^+ + e^- \Rightarrow Ag$	+0.80

(i) Draw a labelled diagram of the cell you could set up in a laboratory to measure the  $E_{\text{cell}}^{\bullet}$  of the reaction involved in cleaning tarnished silver.



(ii) When [Al<sup>3+</sup>] ions are charged in a half cell the  $E_{cell}$  value for the half cell is given by

$$E_{\text{cell}} = E_{\text{cell}}^{\text{o}} - \frac{RT}{nF} \ln \left[ \text{Al}^{3+} \right]$$

Where R = gas constant

T =temperature in kelvin

 $F = \text{Faraday constant}, 9.65 \times 10^4 \text{ C mol}^{-1}$ 

n = number of electrons transferred

Calculate the electrode potential of a cell at T = 298 K and  $[Al^{3+}] = 0.1$  mol dm<sup>-3</sup>.

Show all your working.

$$E_{\text{cell}} = \dots V$$
 [2]

#### **END OF QUESTION PAPER**

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.