Candidate Name	Centre Number			Candidate Number						



**AS BIOLOGY** 

#### **COMPONENT 2**



**Biodiversity and Physiology Of Body Systems** 

**SPECIMEN PAPER** 

1 hour 30 minutes

For Examiner's use only				
Question	Maximum	Mark		
	Mark	Awarded		
1.	8			
2.	20			
3.	15			
4.	12			
5.	11			
6.	9			
Total	75			

## **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need a calculator and a ruler.

### **INSTRUCTIONS TO CANDIDATES**


Use black ink or black ball-point pen. Do not use gel pen. Do not use correction fluid. Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** guestions.

Write your answers in the spaces provided in this booklet.

#### INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question. The assessment of the quality of extended response (QER) will take place in question 6.

1. Different scientists interpret available scientific evidence in different ways. The models below show possible evolutionary relationships between different species of 'big cat'. Model A is based solely on morphological evidence and model B is based on biochemical analyses.



.....

.....

(c)	Samples of mitochondrial DNA from a lion, a tiger and a snow leopard were
	analysed. The base sequences for the same section of their mitochondrial
	DNA are shown below.

	10	20	30	40
Lion	ATACGTGTAT	ACGTGTACGT	GTGTACGTGT	GTACGTGTGT
Tiger	ACGTGTACGT	GTGTATACGT	GTACGTGTGT	ACGTGTGTAT
Snow Leopard	ATACGTGTAC	GTGTATACGT	GTACGTGTGT	ACGTGTGTAC

Species 1	Species 2	Number of bases in common
Lion	Tiger	20
Tiger	Snow Leopard	31
Lion	Snow Leopard	25

	Based on this information which of the two proposed models is supported by these results? Explain your answer.	[1]
(d)	Human mitochondrial DNA codes for proteins essential for some of the reactions which take place within the mitochondrion. Name <b>two</b> forms of R which are essential for the synthesis of these proteins within the mitochond matrix.	


R

2. Plants need water for transport, support and photosynthesis among other biochemical processes. Water uptake from the soil is largely passive and the flow of water from the roots to the leaves is driven in part by the loss of water through the stomata.

(a)	State the name of the process by which plants lose water from their leaves	<b>S</b> .
		[1]

The apparatus shown in the diagram was used to study how wind speed affects the rate at which a leafy shoot takes up water.

The same shoot was exposed to air moving at different speeds. At each speed the distance moved by the air bubble in 5 minutes was recorded. The results were then used to calculate the rate of water uptake by the leafy shoot in mm<sup>3</sup> min<sup>-1</sup>.



(b) The table below shows the results of the investigation.

Wind speed (m s <sup>-1</sup> )	0	5.0	10.0	15	20	25.0
Distance travelled by air bubble in 5 minutes (mm)	64	98	129	163	175	151
Rate of water uptake (mm³ min⁻¹)	10.0	15.3	20.2	25.5	27.4	

(i)	Using the formula below, calculate the rate of water uptake by	
.,	the plant in mm <sup>3</sup> min <sup>-1</sup> at a wind speed of 25 m s <sup>-1</sup> .	[3]

Volume of water = distance travelled by bubble  $\times \pi \times \text{radius}^2$  ( $\pi = 3.14$ )

	Answer = mm³ r	nin <sup>-</sup>
(ii)	Explain why the rate of water uptake increased as the wind speed increased from 0 to 20 m s <sup>-1</sup> .	[3]

The results show, that at a wind speed above 20 m s<sup>-1</sup>, the rate of water (c) uptake by the leafy shoot decreased. One explanation for this decrease could be due to the closure of the stomata by guard cells. Describe the role of each of the following in the operation of guard cells in closing stomata. (i) K+ (potassium ions). [1] (ii) Water potential of the cells. [1] (iii) Unevenly thickened cell walls. [1] (d) Suggest how you could adapt the experimental technique given to determine the location of the highest density of stomata. Predict the results you would expect. [3]


	AS BIOLOGY Specimen Assessment Mate	eriais 29
Man (i)	ny plants are adapted to living in regions of low water availability.  What term is used to describe this type of plant?	[1]
(1)	what term is used to describe this type of plant:	ניו
 (ii)	The photomicrograph below shows a section through a leaf sho number of adaptations to living in an arid environment.	wing a
ころの一般のできません。		
	ntify three features that can be seen in the photomicrograph and expeach helps the plant to reduce water loss.	plain [6]
Feat	ture 1:	
Feat	ture 2:	

Feature 3:	

20

3. The mammalian gut is divided into several regions, each adapted to carry out different functions. The conditions in each region are different and parasites that inhabit the gut must be able to adapt to these conditions.

The diagram below shows how the pH of the alimentary canal changes in different regions.



(a) Explain the cause of the following:

(i)	the increase in pH in region C;	[1]
(ii)	the decrease in pH in region D.	[2]

(b)	(i)	Using letters from the diagram, identify the regions of the human alimentary canal where you would expect to find the tapeworm <i>Taenia solium</i> in an infected person.	[1]
	(ii)	How is the tapeworm adapted to survive in these regions?	[2]
(c)	which proglo averag	Taenia solium usually have about 1 000 proglottids (segments), eac contains both male and female reproductive systems. A mature of ottid can contain about 50 000 eggs and a mature tapeworm releases of six mature proglottids each day. Each tapeworm can survive for 25 years.	s an
	(i)	Suggest why Taenia solium needs to produce so many eggs.	[1]
	(ii)	Estimate the number of eggs that could be produced by a single <i>Taenia solium</i> during its lifetime. Show your workings and give you answer in standard form to two significant figures.	 [3]
		Answer eggs	3
(d)		n why both humans and the tapeworm <i>Taenia solium</i> are described <b>heterotrophic</b> but only the human has <b>holozoic</b> nutrition.	as [3]

Papua New Guinea. The passage below describes some of the initial observations made.	
"The animal is a mammal approximately 1.2 m long and 0.6 m high. It has sharp upper and lower incisors with well-developed canines; the first premolars are sharp with serrated edges while the rear premolars are similar to the molars and have strong ridges. Total gut length is approximately 15.0m (compared to 30.8 m in a sheep and 5.1 m in a dog of similar size). The ratio of body length to intestine length is 1:13 (compared to 1:25 in a sheep and 1:6 in a dog of similar size)."	
	2]
	••
	••
	•


The body of an animal, previously unknown to science, was discovered in

15

(e)

Insects and mammals have both evolved to live in terrestrial environments. One of the main problems they have had to overcome is the absorption of oxygen from the air. To do this both types of organism have evolved internal gas exchange systems.								
(a) (i)	(i) Name the sites of gas exchange in insects and mammals. [1]							
	Insects							
	Mammals							
(ii)	Explain why insects and mammals have internal gas exchange systems.							
insects and m % oxygen in air	before gas exchange  after gas exchange	Insects 21% 17 – 18%	Mammals 21% 13.5 - 16%					
Surface Area body tissue	of gas exchange surface : Volume of	Very high	High					
Association v	None	Close						
	required	No	Yes					
Haemoglobin	num thickness of body tissue (approximate 20 mm no limi							
	ckness of body tissue (approximate	20 mm	no limit					

The diagram below represents the circulatory system of a fish.



(C)	name the type of circulatory system.	[1]
(d)	Fish have evolved gills that allow gas exchange. Water contains approximately 1% dissolved oxygen but fish are able to absorb about 80% the available oxygen and remain highly active.	of
	Describe how the structure of their gills enables bony fish to be so efficient gas exchange.	at [4]

(a)

5. In recent years scientists have become increasingly concerned about the global decrease in **biodiversity**. A higher than normal rate of **extinction** caused by humans has been proposed as the main reason for this decrease. In the UK, hedges have been identified as being habitats that have a high biodiversity.

State what is meant by the terms:	[2]
biodiversity	
extinction	

(b) To assess the change in biodiversity the presence of some types of plant growing in a 100 m length of hedge was surveyed in 2000 and again in 2010. Each of these species provide a source of food for a wide variety of animals.

A diversity index was calculated based on the results. These are shown in the table below.

Species (description)	Number of plants present per species				
(description)	2000	2010			
Hawthorn dense structure; fairly slow growing; grows to 5 – 14m tall	37	8			
Hazel open structure; grows to 4 – 8 m tall; grows many shoots directly from the roots	21	10			
a climbing plant that covers the ground and grows up other plants to get light	36	41			
Elder fast growing shrub usually between 3 and 8 m tall	12	50			
Bramble produces long stems but usually does not exceed 2 m in height	61	12			
DIVERSITY INDEX		0.60			

(i)	Using the formula and the table given below, calculate the Diversity Index for the year 2000 data.	[3]
Where:	N = total number of individual plants n = number of individuals per species $\Sigma$ = sum of	


Species	n	( <b>n-1</b> )	n(n-1)
Hawthorn	37		
Hazel	21		
lvy	36		
Elder	12		
Bramble	61		
	N =		$\sum n(n-1) =$
	N-1 =		
	N(N-1) =	·	
	Diversity Index	=	
(ii)	maintained as h corresponding of Using the inform what conclusion	edges, but are change in the b nation given ar is can be made	ved each year and many are not being turning into lines of trees with a biodiversity of the habitat. Indeed the values for the Diversity Index, about the change in biodiversity in the nation for this change.

# AS BIOLOGY Specimen Assessment Materials 38

(c)	Describe how a sweep net could be used to estimate the Diversity index of insects at the base of a hedge.	[3]
•••••		••••
	Γ	

[9QER]

Frogs have a larval form called tadpoles. As tadpoles, they obtain oxygen from water which has a relatively low oxygen content, but as adults, oxygen is absorbed from the air in simple lungs. This change is accompanied by changes in the properties of the haemoglobin of the larval and adult frogs, which are similar to that observed in foetal and adult humans. Both larval and adult frog haemoglobin can adjust to changing oxygen demands and availability. The graph below shows the effect of pH on haemoglobin from tadpole and adult frogs.



With reference to the information provided, explain how the properties of haemoglobin enable tadpoles and frogs to gain the oxygen required for survival. (The quality of extended response will be assessed in this question.)


AS BIOLOGY Specimen Assessment Materials 40


AS BIOLOGY Specimen Assessment Materials 41