

|               |               |                  |
|---------------|---------------|------------------|
| Surname       | Centre Number | Candidate Number |
| First name(s) |               | 2                |



**GCE AS**

B410U20-1



**FRIDAY, 27 MAY 2022 – AFTERNOON**

**CHEMISTRY – AS component 2**

**Energy, Rate and Chemistry of Carbon Compounds**

1 hour 30 minutes

**ADDITIONAL MATERIALS**

In addition to this examination paper, you will need a:

- calculator;
- **Data Booklet** supplied by WJEC.

**INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen. Do not use gel pen or correction fluid. You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

**Section A** Answer **all** questions.

**Section B** Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

Candidates are advised to allocate their time appropriately between **Section A (10 marks)** and **Section B (70 marks)**.

**INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

The assessment of the quality of extended response (QER) will take place in **Q.10(a)**.

**Section A**

**Section B**

| For Examiner's use only |              |              |
|-------------------------|--------------|--------------|
| Question                | Maximum Mark | Mark Awarded |
| <b>1. to 7.</b>         | <b>10</b>    |              |
| <b>8.</b>               | <b>13</b>    |              |
| <b>9.</b>               | <b>15</b>    |              |
| <b>10.</b>              | <b>13</b>    |              |
| <b>11.</b>              | <b>19</b>    |              |
| <b>12.</b>              | <b>10</b>    |              |
| <b>Total</b>            | <b>80</b>    |              |

B410U201  
01



JUN22B410U20101

**SECTION A**Answer **all** questions.

1. Name the compound  $(\text{CH}_3)_4\text{C}$ . [1]

.....

2. State the meaning of the term 'heterolytic bond fission'. [1]

.....

.....

3. Explain why propanoic acid is soluble in water but propane is not. [2]

.....

.....

.....

4. Propanoic acid reacts with magnesium to form magnesium propanoate and hydrogen.  
Write an equation for this reaction. [1]

.....




5. The enthalpy change for a reversible reaction is  $-98 \text{ kJ mol}^{-1}$ .

(a) On the axes below draw the energy profile for this reaction.

Label the enthalpy change.

[1]



(b) The activation energy for the backward reaction is  $234 \text{ kJ mol}^{-1}$ .

Calculate the activation energy for the forward reaction.

[1]

Activation energy = .....  $\text{kJ mol}^{-1}$

6. A gas cylinder for a barbecue contains 9.0 kg of propane.

Calculate the number of propane molecules in the cylinder.

[2]

Molecules of propane = .....

7. State how many isomers are represented by the formula  $\text{C}_5\text{H}_{12}$ .

[1]

.....



**SECTION B**Answer **all** questions.

8. Propane and propene are typical examples of hydrocarbons.
- (a) Describe the nature of the bonding in propene and explain how this governs its chemical behaviour.

A diagram may be used in support of your answer. [4]

.....

.....

.....

.....

.....

.....

- (b) Propene can undergo polymerisation to form poly(propene).
- Draw the repeating unit in poly(propene). [1]



(c) Although propane is generally unreactive it can react with chlorine in sunlight to form 1-chloropropane as one of the organic products.

(i) Name the type of reaction mechanism which occurs in this case. [1]

.....

(ii) Write the mechanism for the reaction to form 1-chloropropane.

Include one termination step. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) A hydrocarbon has a relative molecular mass of 136. The percentage composition, by mass, is C 88.1%; H 11.9%.

Calculate both the empirical and molecular formulae of the compound.

You **must** show your working. [3]

Empirical formula .....

Molecular formula .....



9. (a) Ethanol can be produced industrially by the hydration of ethene.



- (i) Name the catalyst used in this production. [1]

.....

- (ii) Calculate the average bond enthalpy for the C — C bond using the enthalpy change of reaction above and the average bond enthalpy values given in the table. [3]

| Bond  | Average bond enthalpy / $\text{kJ mol}^{-1}$ |
|-------|--|
| C = C | 612  |
| C — H | 413  |
| C — O | 360  |
| O — H | 463  |

Average bond enthalpy of C — C = .....  $\text{kJ mol}^{-1}$



- (b) (i) State the meaning of the term 'standard enthalpy change of combustion',  $\Delta_c H^\theta$ . [2]

.....

.....

.....

.....

- (ii) The enthalpy change of combustion of ethanol is  $-1370 \text{ kJ mol}^{-1}$ .

The density of ethanol is  $0.789 \text{ g cm}^{-3}$ .

Calculate the heat energy released, in kJ, when  $0.350 \text{ dm}^3$  of ethanol is burned.

Give your answer to an **appropriate** number of significant figures. [3]

Heat energy released = ..... kJ



(c) Ethanol can be heated under reflux with propanoic acid in the presence of concentrated sulfuric acid to form an ester.

(i) Draw a labelled diagram of the apparatus you could use for heating under reflux. [3]

(ii) Explain how this apparatus prevents escape of vapour and give a reason why the escape of vapour should be prevented. [2]

.....

.....

.....

.....

(iii) Draw the structure of the ester that forms. [1]





**BLANK PAGE**


**PLEASE DO NOT WRITE  
ON THIS PAGE**

B410U201  
09





- (b) 1-Chlorobutane can be converted into 2-iodobutane in a two-stage synthesis.



Stage 1 has a 25% yield and stage 2 has a 92% yield.

Calculate the mass of 2-iodobutane made from 37.6 g of 1-chlorobutane.

[3]

Mass of 2-iodobutane = ..... g

- (c) Chlorofluorocarbons, CFCs, were used for a variety of purposes but have now been replaced by hydrofluorocarbons, HFCs.

Explain why HFCs have replaced CFCs.

[4]

.....

.....

.....

.....

.....

.....


13



11. (a) A student carried out an experiment to study the rate of the reaction between calcium carbonate and hydrochloric acid.



He used the following apparatus to measure the total mass of the reagents and the flask every 30 s for 6 minutes.



The solution remained at room temperature and the reaction was still in progress when the final measurement was taken.

His results are shown below.

| Time / min | Mass of reagents + flask / g |
|------------|------------------------------|
| 0          | 171.50                       |
| 0.5        | 171.37                       |
| 1.0        | 171.29                       |
| 1.5        | 171.23                       |
| 2.0        | 171.19                       |
| 2.5        | 171.12                       |
| 3.0        | 171.07                       |
| 3.5        | 171.02                       |
| 4.0        | 170.98                       |
| 4.5        | 170.94                       |
| 5.0        | 170.91                       |
| 5.5        | 170.89                       |
| 6.0        | 170.87                       |



- (i) Suggest why cotton wool was placed in the neck of the flask. [1]

.....


.....

- (ii) Briefly describe a different experimental method, other than loss of mass, that would allow the rate of this reaction to be determined. [2]

.....

.....

- (iii) Complete the plot for the results of the experiment and draw a line of best fit. [3]



- (iv) Use the graph to calculate the rate of reaction, in grams per minute, at 1 minute.

[2]

Rate = .....  $\text{g min}^{-1}$ 

- (v) He used 1.50 g of calcium carbonate and  $40.0 \text{ cm}^3$  of  $1.50 \text{ mol dm}^{-3}$  hydrochloric acid.

Calcium carbonate is the limiting reactant. Calculate the mass of carbon dioxide that would have been lost if the reaction had been allowed to go to completion. [2]

Mass of carbon dioxide = ..... g

- (vi) He then repeated the experiment using 1.50 g of powdered calcium carbonate.

Sketch on the graph in part (iii) the curve he would expect to obtain. Explain any differences in the curves. [3]

.....

.....

.....

.....

.....

.....



- (b) Another student carried out an experiment to study the enthalpy change for the reaction between calcium carbonate and hydrochloric acid.

She reacted 2.50 g of the carbonate with 50.0 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> hydrochloric acid in a polystyrene cup. The acid was in excess.

She used a thermometer that was accurate to  $\pm 0.1$  °C and the temperature rose from 19.2 °C to 21.3 °C.

- (i) Calculate the molar enthalpy change for this reaction, in kJ mol<sup>-1</sup>. [3]

$$\Delta H^\theta = \dots\dots\dots \text{kJ mol}^{-1}$$

- (ii) Calculate the percentage error in the temperature rise recorded. [1]

$$\text{Percentage error} = \dots\dots\dots \%$$

- (iii) She repeated the experiment but used 25.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> hydrochloric acid.

Predict the temperature change in this reaction. Give a reason for your answer. [1]

.....  
 .....


- (iv) She repeated the experiment again but used 50.0 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> nitric acid.

Predict the temperature change in this reaction. Give a reason for your answer. [1]

.....  
 .....



12. Study the reaction scheme shown below and the other information that follows.



Compound **L** is a hydrocarbon. It does not show *E-Z* isomerism and its mass spectrum shows a molecular ion peak at  $m/z$  56.

The  $^1\text{H}$ NMR spectrum for compound **K** shows 3 peaks and the ratio of the peak areas is 6:1:2.

The  $^{13}\text{C}$  NMR spectrum for compound **N** shows 3 peaks.

Compound **O** does not react with sodium carbonate.





(a) Identify compounds **K**, **L**, **M** and **N**. Give your reasoning.

[8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Name the homologous series to which compound **O** belongs.

[1]

.....

(c) State the reagent(s) and conditions needed for the conversion of compound **K** to compound **L**.

[1]

.....

**END OF PAPER**

10







**BLANK PAGE**

**PLEASE DO NOT WRITE  
ON THIS PAGE**





GCE AS

B410U20-1A



FRIDAY, 27 MAY 2022 – AFTERNOON

**CHEMISTRY – AS component 2**  
**Data Booklet**

|                                     |  |
|-------------------------------------|--|
| Avogadro constant                   | $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$               |
| molar gas constant                  | $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$               |
| molar gas volume at 273 K and 1 atm | $V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$                 |
| molar gas volume at 298 K and 1 atm | $V_m = 24.5 \text{ dm}^3 \text{ mol}^{-1}$                 |
| Planck constant                     | $h = 6.63 \times 10^{-34} \text{ Js}$                      |
| speed of light                      | $c = 3.00 \times 10^8 \text{ ms}^{-1}$                     |
| density of water                    | $d = 1.00 \text{ g cm}^{-3}$                               |
| specific heat capacity of water     | $c = 4.18 \text{ J g}^{-1} \text{ K}^{-1}$                 |
| ionic product of water at 298 K     | $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ |
| fundamental electronic charge       | $e = 1.60 \times 10^{-19} \text{ C}$                       |

temperature (K) = temperature (°C) + 273

$1 \text{ dm}^3 = 1000 \text{ cm}^3$   
 $1 \text{ m}^3 = 1000 \text{ dm}^3$   
1 tonne = 1000 kg  
 $1 \text{ atm} = 1.01 \times 10^5 \text{ Pa}$

| Multiple  | Prefix | Symbol |
|-----------|--------|--------|
| $10^{-9}$ | nano   | n      |
| $10^{-6}$ | micro  | $\mu$  |
| $10^{-3}$ | milli  | m      |



| Multiple | Prefix | Symbol |
|----------|--------|--------|
| $10^3$   | kilo   | k      |
| $10^6$   | mega   | M      |
| $10^9$   | giga   | G      |

## Infrared absorption values





| Bond                     | Wavenumber / $\text{cm}^{-1}$ |
|--------------------------|-------------------------------|
| C — Br                   | 500 to 600                    |
| C — Cl                   | 650 to 800                    |
| C — O                    | 1000 to 1300                  |
| C = C                    | 1620 to 1670                  |
| C = O                    | 1650 to 1750                  |
| C $\equiv$ N             | 2100 to 2250                  |
| C — H                    | 2800 to 3100                  |
| O — H (carboxylic acid)  | 2500 to 3200 (very broad)     |
| O — H (alcohol / phenol) | 3200 to 3550 (broad)          |
| N — H                    | 3300 to 3500                  |

<sup>13</sup>C NMR chemical shifts relative to TMS = 0

| Type of carbon | Chemical shift, $\delta$ (ppm) |
|----------------|--------------------------------|
|----------------|--------------------------------|



**$^1\text{H}$  NMR chemical shifts relative to TMS = 0**

| Type of proton  | Chemical shift, $\delta$ (ppm) |
|---|--------------------------------|
| $-\text{CH}_3$  | 0.1 to 2.0                     |
| $\text{R}-\text{CH}_3$  | 0.9                            |
| $\text{R}-\text{CH}_2-\text{R}$   | 1.3                            |
| $\text{CH}_3-\text{C}\equiv\text{N}$  | 2.0                            |
| $\text{CH}_3-\text{C}(=\text{O})$   | 2.0 to 2.5                     |
| $-\text{CH}_2-\text{C}(=\text{O})$  | 2.0 to 3.0                     |
|    | 2.2 to 2.3                     |
| $\text{HC}-\text{Cl}$ or $\text{HC}-\text{Br}$                                      | 3.1 to 4.3                     |
| $\text{HC}-\text{O}$  | 3.3 to 4.3                     |
| $\text{R}-\text{OH}$  | 4.5 *                          |
| $-\text{C}=\text{CH}$   | 4.5 to 6.3                     |
| $-\text{C}=\text{CH}-\text{CO}$   | 5.8 to 6.5                     |
|  | 6.5 to 7.5                     |
|  | 6.5 to 8.0                     |
|  | 7.0 *                          |
| $\text{R}-\text{C}(=\text{O})\text{H}$  | 9.8 *                          |
| $\text{R}-\text{C}(=\text{O})\text{OH}$   | 11.0 *                         |

\*variable figure dependent on concentration and solvent

# THE PERIODIC TABLE

Group 1 2 3 4 5 6 7 0

Period 1 2 3 4 5 6 7

| Period | 1                             | 2                             | p block                       |                               |                                |                                |                                  |                                   |                               |                                  |                                   |                               |                                  |                                   |                                |                                  |                               |
|--------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|----------------------------------|-----------------------------------|-------------------------------|----------------------------------|-----------------------------------|-------------------------------|----------------------------------|-----------------------------------|--------------------------------|----------------------------------|-------------------------------|
| 1      | 1.01<br>H<br>Hydrogen<br>1    |                               |                               |                               |                                |                                |                                  |                                   |                               |                                  |                                   |                               | 4.00<br>He<br>Helium<br>2        |                                   |                                |                                  |                               |
| 2      | 6.94<br>Li<br>Lithium<br>3    | 9.01<br>Be<br>Beryllium<br>4  | 10.8<br>B<br>Boron<br>5       | 12.0<br>C<br>Carbon<br>6      | 14.0<br>N<br>Nitrogen<br>7     | 16.0<br>O<br>Oxygen<br>8       | 19.0<br>F<br>Fluorine<br>9       | 20.2<br>Ne<br>Neon<br>10          | 27.0<br>Al<br>Aluminium<br>13 | 28.1<br>Si<br>Silicon<br>14      | 31.0<br>P<br>Phosphorus<br>15     | 32.1<br>S<br>Sulfur<br>16     | 35.5<br>Cl<br>Chlorine<br>17     | 40.0<br>Ar<br>Argon<br>18         |                                |                                  |                               |
| 3      | 23.0<br>Na<br>Sodium<br>11    | 24.3<br>Mg<br>Magnesium<br>12 | 45.0<br>Sc<br>Scandium<br>21  | 50.9<br>V<br>Vanadium<br>23   | 52.0<br>Cr<br>Chromium<br>24   | 54.9<br>Mn<br>Manganese<br>25  | 55.8<br>Fe<br>Iron<br>26         | 58.7<br>Ni<br>Nickel<br>28        | 58.9<br>Co<br>Cobalt<br>27    | 58.9<br>Rh<br>Rhodium<br>45      | 63.5<br>Cu<br>Copper<br>29        | 65.4<br>Zn<br>Zinc<br>30      | 72.6<br>Ge<br>Germanium<br>32    | 74.9<br>As<br>Arsenic<br>33       | 79.0<br>Se<br>Selenium<br>34   | 79.9<br>Br<br>Bromine<br>35      | 83.8<br>Kr<br>Krypton<br>36   |
| 4      | 39.1<br>K<br>Potassium<br>19  | 40.1<br>Ca<br>Calcium<br>20   | 88.9<br>Y<br>Yttrium<br>39    | 92.9<br>Nb<br>Niobium<br>41   | 95.9<br>Mo<br>Molybdenum<br>42 | 98.9<br>Tc<br>Technetium<br>43 | 101<br>Ru<br>Ruthenium<br>44     | 106<br>Pd<br>Palladium<br>46      | 103<br>Rh<br>Rhodium<br>45    | 108<br>Ag<br>Silver<br>47        | 112<br>Cd<br>Cadmium<br>48        | 115<br>In<br>Indium<br>49     | 119<br>Sn<br>Tin<br>50           | 122<br>Sb<br>Antimony<br>51       | 127<br>Te<br>Tellurium<br>52   | 127<br>I<br>Iodine<br>53         | 131<br>Xe<br>Xenon<br>54      |
| 5      | 85.5<br>Rb<br>Rubidium<br>37  | 87.6<br>Sr<br>Strontium<br>38 | 139<br>La<br>Lanthanum<br>57  | 181<br>Ta<br>Tantalum<br>73   | 184<br>W<br>Tungsten<br>74     | 186<br>Re<br>Rhenium<br>75     | 190<br>Os<br>Osmium<br>76        | 195<br>Pt<br>Platinum<br>78       | 192<br>Ir<br>Iridium<br>77    | 197<br>Au<br>Gold<br>79          | 201<br>Hg<br>Mercury<br>80        | 204<br>Tl<br>Thallium<br>81   | 207<br>Pb<br>Lead<br>82          | 209<br>Bi<br>Bismuth<br>83        | (210)<br>Po<br>Polonium<br>84  | (210)<br>At<br>Astatine<br>85    | (222)<br>Rn<br>Radon<br>86    |
| 6      | 133<br>Cs<br>Caesium<br>55    | 137<br>Ba<br>Barium<br>56     | (227)<br>Fr<br>Francium<br>87 | (226)<br>Ra<br>Radium<br>88   | (227)<br>Ac<br>Actinium<br>89  | (253)<br>Fm<br>Fermium<br>100  | (254)<br>Es<br>Einsteinium<br>99 | (256)<br>Md<br>Mendelevium<br>101 | (253)<br>Fm<br>Fermium<br>100 | (254)<br>Es<br>Einsteinium<br>99 | (256)<br>Md<br>Mendelevium<br>101 | (253)<br>Fm<br>Fermium<br>100 | (254)<br>Es<br>Einsteinium<br>99 | (256)<br>Md<br>Mendelevium<br>101 | (254)<br>No<br>Nobelium<br>102 | (257)<br>Lr<br>Lawrencium<br>103 |                               |
| 7      | (223)<br>Fr<br>Francium<br>87 | (226)<br>Ra<br>Radium<br>88   | (227)<br>Ac<br>Actinium<br>89 | (227)<br>Ac<br>Actinium<br>89 | (227)<br>Ac<br>Actinium<br>89  | (227)<br>Ac<br>Actinium<br>89  | (227)<br>Ac<br>Actinium<br>89    | (227)<br>Ac<br>Actinium<br>89     | (227)<br>Ac<br>Actinium<br>89 | (227)<br>Ac<br>Actinium<br>89    | (227)<br>Ac<br>Actinium<br>89     | (227)<br>Ac<br>Actinium<br>89 | (227)<br>Ac<br>Actinium<br>89    | (227)<br>Ac<br>Actinium<br>89     | (227)<br>Ac<br>Actinium<br>89  | (227)<br>Ac<br>Actinium<br>89    | (227)<br>Ac<br>Actinium<br>89 |

**Key**

|      |                      |
|------|----------------------|
| Ar   | Symbol               |
| Name | atomic number        |
| Z    | relative atomic mass |