



---

# **GCE A LEVEL MARKING SCHEME**

---

**AUTUMN 2021**

**A LEVEL  
CHEMISTRY - COMPONENT 3  
A410U30-1**

## **INTRODUCTION**

This marking scheme was used by WJEC for the 2021 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

## COMPONENT 3: CHEMISTRY IN PRACTICE

### AUTUMN 2021 MARK SCHEME

#### GENERAL INSTRUCTIONS

##### Recording of marks

Examiners must mark in red ink.

One tick must equate to one mark, apart from extended response questions where a level of response mark scheme is applied.

Question totals should be written in the box at the end of the question.

Question totals should be entered onto the grid on the front cover and these should be added to give the script total for each candidate.

##### Extended response questions

A level of response mark scheme is applied. The complete response should be read in order to establish the most appropriate band. Award the higher mark if there is a good match with content and communication criteria. Award the lower mark if either content or communication barely meets the criteria.

##### Marking rules

All work should be seen to have been marked.

Marking schemes will indicate when explicit working is deemed to be a necessary part of a correct answer.

Crossed out responses not replaced should be marked.

### Marking abbreviations

The following may be used in marking schemes or in the marking of scripts to indicate reasons for the marks awarded.

cao = correct answer only  
ecf = error carried forward  
bod = benefit of doubt

Credit should be awarded for correct and relevant alternative responses which are not recorded in the mark scheme.

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
1	(a)	(i)	$\text{C}_3\text{H}_8\text{O(l)} + 4\frac{1}{2}\text{O}_2\text{(g)} \rightarrow 3\text{CO}_2\text{(g)} + 4\text{H}_2\text{O(l)}$ balanced equation (1) state symbols (1)	2			2		
		(ii)	spirit burner containing propan-1-ol and metal can containing water accept beaker in place of metal can (1)  known volume / mass of water added to metal can (1)  award (1) for <b>both</b> measurements <ul style="list-style-type: none"> <li>initial and final temperature of the water (allow temperature change of the water)</li> <li>mass of the burner before and after heating (allow change in mass of the burner)</li> </ul> use of $\Delta H = -\frac{m \times c \times \Delta T}{n}$ (1)	2			4		4
	(b)	(i)	suitable scale selected on <i>x</i> -axis and <i>y</i> -axis (1)  all points plotted correctly (tolerance $\pm\frac{1}{2}$ square) and line of best fit drawn (1)		2		2	2	
		(ii)	as the number of moles of ethanol (in the mixture) increases the heat evolved (in kJ) decreases  linear relationship – neutral answer			1	1	1	

Question				Marking details	Marks available					
					AO1	AO2	AO3	Total	Maths	Prac
		(iii)		-2005 accept 2005		1		1		
		(iv)		n(CH <sub>3</sub> CH <sub>2</sub> OH) from graph = 0.0072 (1) mass of CH <sub>3</sub> CH <sub>2</sub> OH = 0.0072 × 46.06 = 0.332 (1)		1	1	2	1	
		(v)		line starting from same point on right-hand side as plotted line (1) gradient greater than plotted line (1)			2	2	2	
				<b>Question 1 total</b>	<b>4</b>	<b>6</b>	<b>4</b>	<b>14</b>	<b>6</b>	<b>4</b>

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
2	(a)		award (1) for either of following <ul style="list-style-type: none"> <li>reacts with phenol to produce the (nucleophilic) phenoxide ion</li> <li>removes the proton from phenol</li> </ul>		1		1		1
	(b)	(i)	benzoic anhydride (accept benzoic acid)		1		1		1
		(ii)	award (1) for either of following  benzoyl chloride more reactive not an equilibrium process			1	1		1
	(c)		dissolve solid in minimum volume of hot ethanol (1)  award (1) each for any <b>two</b> of following <ul style="list-style-type: none"> <li>heat ethanol using a hot plate / hot water bath / electric heating mantle</li> <li>filter off any insoluble impurities whilst hot</li> <li>allow to cool (to crystallise solid)</li> <li>filter and dry</li> </ul>		3		3		3

Question		Marking details		Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
	(d)		$n(\text{C}_6\text{H}_5\text{OH}) = \frac{1.5}{94.06} = 0.016 \quad (1)$ mass of $\text{C}_7\text{H}_5\text{OCl} = 3.0 \times 1.2 = 3.6 \text{ g}$ $n(\text{C}_7\text{H}_5\text{OCl}) = \frac{3.6}{140.55} = 0.026 \quad \Rightarrow \text{in excess } (1)$ theoretical mass of $(\text{C}_{13}\text{H}_{10}\text{O}_2) = 0.016 \times 198.1 = 3.2 \text{ g} \quad (1)$ percentage yield = $\frac{2.9}{3.2} \times 100 = 91 \quad (1)$		3	1	4	2	
	(e)		award (1) each for any <b>two</b> of following  NMR HPLC GC mass spectra melting temperature		2		2		2
	(f)	(i)	$\text{CHI}_3 \quad (1)$  $\begin{array}{c} \text{CH}_3 \\   \\ \text{R}-\text{C}-\text{OH} \\   \\ \text{H} \end{array} \quad (1)$		2		2		2
		(ii)	acidified potassium dichromate solution $(1)$  colour change from orange to green $(1)$	2			2		2



Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
		(iii)	aldehyde		1		1		1
		(iv)	award (2) for <b>all</b> four ester structures award (1) for any <b>two</b> ester structures  <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <math display="block">\begin{array}{c} \text{O} \\ \parallel \\ \text{H}_3\text{C}-\text{CH}_2-\text{C} \\   \\ \text{O}-\text{CH}_3 \end{array}</math> </div> <div style="text-align: center;"> <math display="block">\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{C} \\   \\ \text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_3 \end{array}</math> </div> </div> <div style="display: flex; justify-content: space-around; align-items: center; margin-top: 20px;"> <div style="text-align: center;"> <math display="block">\begin{array}{c} \text{O} \\ \parallel \\ \text{H}_3\text{C}-\text{C} \\   \\ \text{O}-\text{CH}_2-\text{CH}_3 \end{array}</math> <p><b>A</b></p> </div> <div style="text-align: center;"> <math display="block">\begin{array}{c} \text{O} \\ \parallel \\ \text{H}-\text{C} \\   \\ \text{O}-\text{CH}-\text{CH}_3 \\   \\ \text{CH}_3 \end{array}</math> <p><b>B</b></p> </div> </div> award (1) for each of <b>A</b> and <b>B</b> correctly identified			4	4		2
<b>Question 2 total</b>				<b>2</b>	<b>13</b>	<b>6</b>	<b>21</b>	<b>2</b>	<b>15</b>

Question	Marking details	Marks available					
		AO1	AO2	AO3	Total	Maths	Prac
3	<p><b>Indicative content</b></p> <ol style="list-style-type: none"> <li><b>Fe(NO<sub>3</sub>)<sub>2</sub></b> identified from <b>pale green</b> solution</li> <li>Fe(NO<sub>3</sub>)<sub>2</sub> added to other five solutions</li> <li><b>Pale green precipitate</b> formed identifies <b>NaOH</b></li> <li>NaOH added to remaining four solutions <ul style="list-style-type: none"> <li><b>two solutions</b> give <b>white precipitate</b> which are <b>soluble in excess</b> NaOH indicating <b>Pb(NO<sub>3</sub>)<sub>2</sub></b> and <b>Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub></b></li> <li><b>two solutions</b> give <b>no observable reaction</b> indicating <b>Ba(NO<sub>3</sub>)<sub>2</sub></b> and <b>KI</b></li> </ul> </li> <li>Take one of the solutions which gave a white precipitate with NaOH and add in turn to both solutions that gave no observable reaction with NaOH <ul style="list-style-type: none"> <li>If white precipitate formed then <b>Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub></b> and <b>Ba (NO<sub>3</sub>)<sub>2</sub></b> both identified</li> <li>If (bright) yellow precipitate formed then <b>Pb(NO<sub>3</sub>)<sub>2</sub></b> and <b>KI</b> both identified</li> </ul> </li> </ol>	2		4	6		4
	<p><b>5-6 marks</b> Method outlined leads to the identification of all six solutions <i>The candidate constructs a relevant, coherent and logically structured account including all key elements of the indicative content. A sustained and substantiated line of reasoning is evident and scientific conventions and vocabulary is used accurately throughout.</i></p> <p><b>3-4 marks</b> Method outlined leads to the identification of Fe(NO<sub>3</sub>)<sub>2</sub> and NaOH and solutions containing amphoteric and non-amphoteric metals <i>The candidate constructs a coherent account including many of the key elements of the indicative content. Some reasoning is evident in the linking of key points and use of scientific conventions and vocabulary is generally sound.</i></p>						

Question				Marking details	Marks available						
					AO1	AO2	AO3	Total	Maths	Prac	
				<p><b>1-2 marks</b>  Method leads to the identification of <math>\text{Fe}(\text{NO}_3)_2</math> and <math>\text{NaOH}</math>  <i>The candidate attempts to link at least two relevant points from the indicative material. Coherence is limited by omission and/or inclusion of irrelevant materials. There is some evidence of appropriate use of scientific conventions and vocabulary.</i></p> <p><b>0 marks</b>  <i>The candidate does not make any attempt or give an answer worthy of credit.</i></p>							
				<b>Question 3 total</b>	<b>2</b>	<b>0</b>	<b>4</b>	<b>6</b>	<b>0</b>	<b>4</b>	

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
4	(a)	(i)	award (3) for <b>all</b> five points award (2) for point 4 <u>and</u> any <b>three</b> others award (1) for any three points  1. Record the mass of an empty crucible + lid 2. Add known mass of CaSO <sub>4</sub> .2H <sub>2</sub> O and replace the crucible lid 3. Heat the crucible (with lid) containing the solid above a Bunsen burner flame / carefully lift lid but do not remove completely 4. Heat to constant mass 5. Weigh crucible + lid + solid (after cooling)		3		3		3
		(ii)	$n(\text{CaSO}_4 \cdot 2\text{H}_2\text{O}) \text{ used} = \frac{5.20}{172.24} = 0.0302 \quad (1)$  mass of anhydrous CaSO <sub>4</sub> formed must be measured  $\text{mass} = 136.2 \times 0.0302 = 4.11 \text{ g} \quad (1)$  accept mass of water lost as an alternative  $\text{mass} = 0.0302 \times 2 \times 18.02 = 1.09 \text{ g}$		2		2	2	
	(b)		$n(\text{CaSO}_4 \cdot 2\text{H}_2\text{O}) = \frac{2.49}{172.24} = 0.0145 \text{ mol} \quad (1)$  $M_r(\text{M}_2\text{SO}_4) = \frac{1.59}{0.0145} = 109.90 \quad (1)$  $M_r(\text{M}) = \frac{(109.90 - 96.1)}{2} = 6.91 \Rightarrow \text{M is lithium} \quad (1)$		2		3	2	
<b>Question 4 total</b>				<b>0</b>	<b>7</b>	<b>1</b>	<b>8</b>	<b>4</b>	<b>3</b>

Question			Marking details	Marks available					
				AO1	AO2	AO3	Total	Maths	Prac
5	(a)		$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$		1		1		
	(b)		$\text{PbO}_2 + 4\text{H}^+ + 2\text{e}^- \rightarrow \text{Pb}^{2+} + 2\text{H}_2\text{O}$			1	1		
	(c)		starch indicator blue/black to colourless	1			1		1
	(d)	(i)	$n(\text{S}_2\text{O}_3^{2-}) = 0.0510 \times \frac{21.95}{1000} = 0.00112$ $n(\text{I}_2) \text{ in } 25.0 \text{ cm}^3 = 0.000560$ $n(\text{I}_2) \text{ formed in step 1} = 8 \times 0.000560 = 0.00448$	1			1	1	
		(ii)	$n(\text{PbO}_2) = 0.00448 \text{ in } 200 \text{ cm}^3 \text{ of solution} \quad (1)$ mass of $\text{PbO}_2 = 0.00448 \times 239 = 1.071 \text{ g}$ percentage $\text{PbO}_2 = \frac{1.071}{1.18} \times 100 = 90.7 \quad (1)$			1	2	2	
	(e)		$\frac{2 \times 0.005}{1.18} \times 100 = 0.85$		1		1	1	
	(f)	(i)	$2\text{PbO}_2(\text{s}) \rightarrow 2\text{PbO}(\text{s}) + \text{O}_2(\text{g})$ ignore state symbols		1		1		

Question			Marking details	Marks available						
				AO1	AO2	AO3	Total	Maths	Prac	
		(ii)	award (1) for conversion / correct units for p, V and T  $n(\text{O}_2) = \frac{pV}{RT} = \frac{1.01 \times 10^5 \times 1.23 \times 10^{-4}}{8.31 \times 873} = 0.00171 \quad (1)$  $n(\text{PbO}_2) = 0.00342$  $\text{mass of PbO}_2 = 0.00342 \times 239 = 0.819 \quad (1)$		3		3	3		
			<b>Question 5 total</b>	<b>3</b>	<b>6</b>	<b>2</b>	<b>11</b>	<b>7</b>	<b>1</b>	

**COMPONENT 3: CHEMISTRY IN PRACTICE**  
**SUMMARY OF MARKS ALLOCATED TO ASSESSMENT OBJECTIVES**

<b>Question</b>	<b>AO1</b>	<b>AO2</b>	<b>AO3</b>	<b>Total</b>	<b>Maths</b>	<b>Prac</b>
<b>1</b>	<b>4</b>	<b>6</b>	<b>4</b>	<b>14</b>	<b>6</b>	<b>4</b>
<b>2</b>	<b>2</b>	<b>13</b>	<b>6</b>	<b>21</b>	<b>2</b>	<b>15</b>
<b>3</b>	<b>2</b>	<b>0</b>	<b>4</b>	<b>6</b>	<b>0</b>	<b>4</b>
<b>4</b>	<b>0</b>	<b>7</b>	<b>1</b>	<b>8</b>	<b>4</b>	<b>3</b>
<b>5</b>	<b>3</b>	<b>6</b>	<b>2</b>	<b>11</b>	<b>7</b>	<b>1</b>
<b>Totals</b>	<b>11</b>	<b>32</b>	<b>17</b>	<b>60</b>	<b>19</b>	<b>27</b>