

## Friday 23 June 2023 – Morning

### A Level Chemistry A

H432/03 Unified chemistry

Time allowed: 1 hour 30 minutes



**You must have:**

- the Data Sheet for Chemistry A

**You can use:**

- a scientific or graphical calculator
- an HB pencil



Please write clearly in black ink. **Do not write in the barcodes.**

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

First name(s)

---

Last name

---

### INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

### INFORMATION

- The total mark for this paper is **70**.
- The marks for each question are shown in brackets [ ].
- Quality of extended response will be assessed in questions marked with an asterisk (\*).
- This document has **24** pages.

### ADVICE

- Read each question carefully before you start your answer.

1 This question is about the first 36 elements in the periodic table:

H																	He
Li	Be											B	C	N	O	F	Ne
Na	Mg											Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr

From these 36 elements **only**, write the symbol for the element(s) that fit each description.

- (a) The **three** elements that, when in a molecule, can form hydrogen bonds with other suitable molecules.

..... [1]

- (b) The element in Period 2 with the successive ionisation energies shown below.

Ionisation number	1st	2nd	3rd	4th	5th	6th	7th	8th
Ionisation energy /kJ mol <sup>-1</sup>	1000	2251	3361	4564	7012	8496	27 107	31 671

..... [1]

- (c) An element that is a solid at RTP with a simple molecular lattice structure.

..... [1]

(d) The **two** elements with atoms containing five unpaired electrons.

..... [2]

(e) The element in Period 3 that exists in the solid state as a network of atoms bonded by strong covalent bonds.

..... [1]

(f) The p-block element in Period 3 that forms a compound with fluorine with octahedral molecules.

..... [1]

(g) The element that forms 1– ions most readily.

..... [1]

(h) The element with an average atomic mass of  $1.244 \times 10^{-22}$  g.

..... [1]

2 These questions are from different areas of chemistry.

(a) This question is about two salts of rubidium (atomic number 37):  $\text{RbClO}_3$  and  $\text{RbClO}_4$ .

(i) The oxidation number of chlorine is different in the two rubidium salts,  $\text{RbClO}_3$  and  $\text{RbClO}_4$ .

What is the name of  $\text{RbClO}_4$ ?

..... [1]

(ii) A student carries out an experiment to determine the enthalpy change of solution of  $\text{RbClO}_3$  using the method below.

- A 2.00 g sample of solid  $\text{RbClO}_3$  is added to water in a well-insulated container. The initial temperature is  $23.0^\circ\text{C}$ .
- The mixture is stirred until all the  $\text{RbClO}_3$  has dissolved. The final temperature is  $21.5^\circ\text{C}$ . The final solution has a mass of 102 g.

Determine the enthalpy change of solution,  $\Delta_{\text{sol}}H$ , of  $\text{RbClO}_3$  in  $\text{kJ mol}^{-1}$ .

Assume that the specific heat capacity of the solution is the same as that of pure water.

$\Delta_{\text{sol}}H(\text{RbClO}_3) = \dots\dots\dots \text{kJ mol}^{-1}$  [3]


(b) A student investigates the rate of a reaction that is 1st order with respect to hydrochloric acid,  $\text{HCl}(\text{aq})$ .

- The student carries out a reaction using  $0.680 \text{ mol dm}^{-3} \text{ HCl}(\text{aq})$ . The initial rate is  $9.52 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$ .
- The student dilutes a different sample of  $0.680 \text{ mol dm}^{-3} \text{ HCl}(\text{aq})$  with water. The pH of this diluted acid is 1.50.
- The student repeats the reaction using the same volume of this diluted acid.

Determine the initial rate using this diluted acid.

initial rate = .....  $\text{mol dm}^{-3} \text{ s}^{-1}$  [3]

(c) The benzenediazonium ion, shown below, is stable at temperatures below 10 °C.



**Benzenediazonium**

Above 10 °C, the benzenediazonium ion reacts with water to form phenol.


The reaction proceeds in a three-step mechanism.

**Step 1** Elimination of nitrogen gas to form a carbocation.

**Step 2** Nucleophilic attack by water.

**Step 3** Proton loss to form the organic product.

Complete the boxes below with intermediates and curly arrows to show the mechanism for this reaction.




[4]

**7**  
**BLANK PAGE**

**DO NOT WRITE ON THIS PAGE**  
**Turn over for the next question**

- 3\* A chemist is investigating compound **A**, shown below, as a potential organic intermediate.



**Compound A**

Describe the type of stereoisomerism shown by compound **A** and suggest **three** reactions of compound **A**, one for each of the three functional groups using reagents of your choice.

In your answer, show stereoisomers of compound **A**, your chosen reactants and conditions, and the structures for the organic products produced.

Mechanisms and equations are **not** required.

**[6]**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional answer space if required.

.....

.....

.....

.....

.....

4 Tutton's salts are 'double salts' with the formula  $X_2Y(ZO_4)_2 \cdot 6H_2O$ .

A Tutton's salt contains two cations:  $X^+$  and  $Y^{2+}$ .

- $X^+$  can be an ion of the Group 1 elements K, Rb, Cs or Fr, or an ammonium ion.
- $Y^{2+}$  can be a 2+ ion of magnesium or an ion of most of the transition elements in Period 4.
- $Z$  can be S or Cr.

$(NH_4)_2Cu(SO_4)_2 \cdot 6H_2O$  is an example of a Tutton's salt.

(a) Predict the formula of a Tutton's salt containing different ions from  $(NH_4)_2Cu(SO_4)_2 \cdot 6H_2O$ .

..... [1]

(b) A student prepares a sample of the Tutton's salt,  $(NH_4)_2Cu(SO_4)_2 \cdot 6H_2O$  using the method shown below.

**Step 1** Dissolve 0.025 mol of ammonium sulfate and 0.025 mol of hydrated copper(II) sulfate,  $CuSO_4 \cdot 5H_2O$ , in water and make up to 50 cm<sup>3</sup>.

**Step 2** Boil the resulting mixture for 2 minutes and allow to cool.

**Step 3** Allow the solvent to evaporate slowly. Pale blue crystals of the Tutton's salt form.

(i) What masses are needed of ammonium sulfate and  $CuSO_4 \cdot 5H_2O$ ?

mass of ammonium sulfate ..... g

mass of  $CuSO_4 \cdot 5H_2O$  ..... g  
[2]

(ii) In **Step 3**, why does the student allow the solvent to evaporate and **not** boil off all the solvent in **Step 2**?

.....  
.....  
..... [1]

- (c) The student dissolves their Tutton's salt in water. A pale blue solution forms.

The student carries out two tests on this aqueous solution.

- (i) The student adds an excess of aqueous ammonia to their aqueous solution of Tutton's salt. A deep blue solution forms.

The complex ion responsible for the deep blue solution has a molar mass of  $167.5 \text{ g mol}^{-1}$ .

Suggest the formula of this complex ion.

..... [1]

- (ii) The student adds  $\text{NaOH(aq)}$  to the aqueous solution of Tutton's salt and warms the mixture.

A precipitate and a gas are formed.

Write the formulae of the precipitate and gas and suggest a test that could confirm the identity of the gas.

Formula of precipitate .....

Formula of gas .....

Test to confirm the identity of the gas .....

.....

..... [3]

- (iii) How could the student carry out a test-tube test to confirm the anion in the Tutton's salt?

.....


.....

.....

..... [2]

- 5 Aspirin tablets are used for pain relief.

The structure of aspirin is shown below.



**Aspirin**

- (a) A student uses the reaction of aspirin with cold NaOH(aq) to determine the mass of aspirin in **one** tablet.

In this reaction, 1 mol of aspirin reacts with 1 mol of cold NaOH(aq).


The student's method is outlined below.

**Step 1** The student reacts **three** aspirin tablets with 100 cm<sup>3</sup> of 0.500 mol dm<sup>-3</sup> NaOH(aq). The NaOH is in excess. A colourless solution forms.

**Step 2** The colourless solution from **Step 1** is made up to 250.0 cm<sup>3</sup> with distilled water.

**Step 3** A 25.00 cm<sup>3</sup> sample of the diluted solution from **Step 2** is titrated with 0.200 mol dm<sup>-3</sup> HCl(aq) in the burette.

The HCl(aq) reacts with excess NaOH(aq) that remains in **Step 1**:



The student repeats the titration to obtain concordant (consistent) titres.

#### Titration results

The trial titre has been omitted.

The burette readings have been read to the nearest 0.05 cm<sup>3</sup>.

	<b>1</b>	<b>2</b>	<b>3</b>
<b>Final reading / cm<sup>3</sup></b>	23.10	45.40	27.40
<b>Initial reading / cm<sup>3</sup></b>	0.00	23.10	5.00

**Analysis of results**

From the results, the student can determine the following.

1. The amount, in mol, of excess NaOH(aq) that remains after the reaction of aspirin with NaOH(aq).
2. The amount, in mol, of NaOH(aq) that reacted with the aspirin.

Use the results to determine the mass, in mg, of aspirin in **one** aspirin tablet.

mass of aspirin in **one** tablet = ..... mg [6]

(b) Aspirin reacts with hot NaOH(aq), under reflux.


(i) Draw a labelled diagram of suitable apparatus for reflux.

[2]

(ii) In this reaction, 1 mol of aspirin reacts with 3 mol of hot NaOH(aq).

Complete the equation for the reaction of aspirin with an excess of hot NaOH(aq).

Show structures for organic compounds.



[3]


16  
BLANK PAGE

DO NOT WRITE ON THIS PAGE



6 This question is about nitric acid, hydrochloric acid and sulfuric acid.

- (a) Nitric acid has 2 single covalent bonds, 1 double covalent bond and 1 dative covalent bond as shown below.



**Nitric acid**

Predict the H–O–N and O–N–O bond angles in nitric acid.

Explain your reasoning.

.....

.....

.....

.....

.....

.....

..... [4]

- (b) Dilute nitric acid reacts with aluminium oxide to form a solution of aluminium nitrate.

(i) Write an equation for this reaction.

..... [2]

(ii) The solution contains nitrate ions,  $\text{NO}_3^-$ .

Draw a 'dot-and-cross' diagram for the  $\text{NO}_3^-$  ion.


Use a different symbol for the extra electron.

[2]

- (c) A mixture of concentrated nitric and hydrochloric acid is called 'aqua regia'. Aqua regia can dissolve gold.

The reaction of aqua regia with gold is a redox reaction which forms chlorauric acid,  $\text{HAuCl}_4$ .

- (i) Balance the half-equation for the oxidation process in this reaction.



[1]

- (ii) In the reduction process in this reaction,  $\text{HNO}_3$  and  $\text{H}^+$  react together to form 2 oxides: **X** ( $M_r = 30$ ) and **Z** ( $M_r = 18$ ).

Determine the formulae of **X** and **Z** and write the half-equation for this reduction.


**X** = .....

**Z** = .....

half-equation ..... [3]

(d) In the UK, most sulfuric acid,  $\text{H}_2\text{SO}_4$ , is manufactured by the Contact process.

One stage in the Contact process involves the equilibrium between sulfur dioxide, oxygen and sulfur trioxide.



This equilibrium is investigated:

**Step 1**  $5.82 \times 10^{-2}$  mol of  $\text{SO}_2$  is mixed with  $7.40 \times 10^{-2}$  mol of  $\text{O}_2$  in a  $2.00 \text{ dm}^3$  container.

**Step 2** The container is sealed and allowed to reach equilibrium at constant temperature.

**Step 3** At equilibrium,  $5.20 \times 10^{-2}$  mol of  $\text{SO}_3$  is formed.

Determine the equilibrium concentrations and calculate  $K_c$ , including units.

$K_c = \dots\dots\dots$  units  $\dots\dots\dots$  [5]

**TURN OVER FOR QUESTION 6 PART (e)**

(e)\* Three reactions involving sulfuric acid are shown below.

**Reaction 1**

Dilute sulfuric acid is reacted with nickel(II) hydroxide to form a green solution.

The solvent is allowed to evaporate leaving hydrated crystals of compound **D**, with the percentage composition by mass: Ni, 22.33%; S, 12.20%; O, 60.87%; H, 4.60%.

**Reaction 2**

Concentrated sulfuric acid is reacted with hydrogen bromide, HBr, to form three products:

- an element which exists as diatomic molecules
- a gaseous compound **E**
- a liquid.

At RTP, 1.00 dm<sup>3</sup> of compound **E** has a mass of 2.67 g.

**Reaction 3**

Concentrated sulfuric acid acts as a catalyst when 2-hydroxypropanoic acid reacts to form compound **F** ( $M_r = 144$ ).

In this reaction, 2 mol of 2-hydroxypropanoic acid forms 1 mol of compound **F** and 2 mol of water.

Identify compounds **D**, **E** and **F** and construct equations for the reactions.

Show structures for any organic compounds.

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Additional answer space if required.

.....

.....

.....

.....

.....

**END OF QUESTION PAPER**

**ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

A large area of lined paper for writing answers. It features a vertical margin line on the left side and horizontal dotted lines for writing. The lines are evenly spaced and extend across the width of the page.

Lined writing area consisting of a vertical solid line on the left and horizontal dotted lines across the page.

A large area of the page is reserved for writing, featuring a vertical solid line on the left side and horizontal dotted lines extending across the page.

---

# OCR

Oxford Cambridge and RSA

## Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.